

Fank1 antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI13291

Product Information

Application WB
Primary Accession Q9DAM9

Other Accession <u>NM_025850</u>, <u>NP_080126</u>

ReactivityHuman, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine **Predicted**Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 38260

Additional Information

Gene ID 66930

Alias Symbol 1700007B22Rik, AI850911

Other Names Fibronectin type 3 and ankyrin repeat domains 1 protein, Germ cell-specific

gene 1 protein, GSG1, Fank1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-Fank1 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions Fank1 antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name Fank1 {ECO:0000312 | MGI:MGI:1914180}

Function Through the activation of JUN and AP-1-mediated transcription, may

regulate apoptosis.

Cellular Location Nucleus. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q8TC84}. Cytoplasm,

cytoskeleton, cilium basal body. Cell projection, cilium. Note=Weakly detected

in the cytoplasm of elongated spermatids.


Tissue Location Mostly restricted to testis (at protein level), including mid to late pachytene

spermatocytes (stages VI-X), diplotene spermatocytes (stage XI), meiotically dividing spermatocytes (stage XII) and spermatids in steps 1-14. Highest levels

References

Hwang K.-C.,et al.Mol. Reprod. Dev. 71:275-283(2005). Carninci P.,et al.Science 309:1559-1563(2005). Zheng Z.,et al.Gene Expr. Patterns 7:777-783(2007).

Images



WB Suggested Anti-Fank1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:100

Positive Control: Mouse Small Intestine

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.