

# OMA1 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI13463

## Product Information

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q96E52</a>
<b>Other Accession</b>	<a href="#">NM_145243</a> , <a href="#">NP_660286</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine, Yeast
<b>Predicted Host</b>	Human, Mouse, Rat, Rabbit, Pig, Chicken, Dog, Guinea Pig, Horse, Bovine
<b>Clonality</b>	Rabbit
<b>Calculated MW</b>	Polyclonal 60120

## Additional Information

<b>Gene ID</b>	115209
<b>Alias Symbol Other Names</b>	2010001O09Rik, DAB1, FLJ33782, MPRP-1, YKR087C, ZMPOMA1 Metalloendopeptidase OMA1, mitochondrial, 3.4.24.-, Metalloprotease-related protein 1, MPRP-1, Overlapping with the m-AAA protease 1 homolog, OMA1, MPRP1
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-OMA1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	OMA1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

<b>Name</b>	OMA1 {ECO:0000303 PubMed:20038677, ECO:0000312 HGNC:HGNC:29661}
<b>Function</b>	Metalloprotease that is part of the quality control system in the inner membrane of mitochondria (PubMed: <a href="#">20038677</a> , PubMed: <a href="#">25605331</a> , PubMed: <a href="#">32132706</a> , PubMed: <a href="#">32132707</a> ). Activated in response to various mitochondrial stress, leading to the proteolytic cleavage of target proteins, such as OPA1, UQCC3 and DELE1 (PubMed: <a href="#">20038677</a> , PubMed: <a href="#">25275009</a> , PubMed: <a href="#">32132706</a> , PubMed: <a href="#">32132707</a> ). Involved in the fusion of the mitochondrial inner membranes by mediating cleavage of OPA1 at S1 position, generating the soluble OPA1 (S-OPA1), which cooperates with the

membrane form (L-OPA1) to coordinate the fusion of mitochondrial inner membranes (PubMed:[31922487](#)). Following stress conditions that induce loss of mitochondrial membrane potential, mediates cleavage of OPA1, leading to excess production of soluble OPA1 (S-OPA1) and negative regulation of mitochondrial fusion (PubMed:[20038677](#), PubMed:[25275009](#)). Involved in mitochondrial safeguard in response to transient mitochondrial membrane depolarization (flickering) by catalyzing cleavage of OPA1, leading to excess production of S-OPA1, preventing mitochondrial hyperfusion (By similarity). Also acts as a regulator of apoptosis: upon BAK and BAX aggregation, mediates cleavage of OPA1, leading to the remodeling of mitochondrial cristae and allowing the release of cytochrome c from mitochondrial cristae (PubMed:[25275009](#)). In depolarized mitochondria, may also act as a backup protease for PINK1 by mediating PINK1 cleavage and promoting its subsequent degradation by the proteasome (PubMed:[30733118](#)). May also cleave UQCRC3 in response to mitochondrial depolarization (PubMed:[25605331](#)). Also acts as an activator of the integrated stress response (ISR): in response to mitochondrial stress, mediates cleavage of DELE1 to generate the processed form of DELE1 (S- DELE1), which translocates to the cytosol and activates EIF2AK1/HRI to trigger the ISR (PubMed:[32132706](#), PubMed:[32132707](#)). Its role in mitochondrial quality control is essential for regulating lipid metabolism as well as to maintain body temperature and energy expenditure under cold-stress conditions (By similarity). Binds cardiolipin, possibly regulating its protein turnover (By similarity). Required for the stability of the respiratory supercomplexes (By similarity).

#### Cellular Location

Mitochondrion inner membrane; Single-pass membrane protein  
{ECO:0000250|UniProtKB:Q9D8H7}


#### Tissue Location

Widely expressed, with strong expression in the heart, skeletal muscle, kidney and liver

## References

Bao Y.-C.,et al.DNA Res. 10:123-128(2003).  
Gregory S.G.,et al.Nature 441:315-321(2006).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Head B.,et al.J. Cell Biol. 187:959-966(2009).

## Images



WB Suggested Anti-OMA1 Antibody Titration: 0.2-1 µg/ml  
Positive Control: 721\_B cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.