

VPS4A antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI13710

Product Information

Application	WB
Primary Accession	Q9UN37
Other Accession	NM_013245 , NP_037377
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48898

Additional Information

Gene ID	27183
Alias Symbol	FLJ22197, SKD1, SKD2, VPS4, VPS4-1, SKD1A
Other Names	Vacuolar protein sorting-associated protein 4A, 3.6.4.6, Protein SKD2, VPS4-1, hVPS4, VPS4A {ECO:0000312 EMBL:AAG01470.1}
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-VPS4A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	VPS4A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	VPS4A {ECO:0000312 EMBL:AAG01470.1}
Function	Involved in late steps of the endosomal multivesicular bodies (MVB) pathway. Recognizes membrane-associated ESCRT-III assemblies and catalyzes their disassembly, possibly in combination with membrane fission. Redistributes the ESCRT-III components to the cytoplasm for further rounds of MVB sorting. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. It is required for proper accomplishment of various processes including the regulation of endosome size, primary cilium organization,

mitotic spindle organization, chromosome segregation, and nuclear envelope sealing and spindle disassembly during anaphase (PubMed:[33186545](#)). Involved in cytokinesis: retained at the midbody by ZFYVE19/ANCHR and CHMP4C until abscission checkpoint signaling is terminated at late cytokinesis. It is then released following dephosphorylation of CHMP4C, leading to abscission (PubMed:[24814515](#)). VPS4A/B are required for the exosomal release of SDCBP, CD63 and syndecan (PubMed:[22660413](#)). Critical for normal erythroblast cytokinesis and correct erythropoiesis (PubMed:[33186543](#)).


Cellular Location

Late endosome membrane {ECO:0000250 | UniProtKB:Q8VEJ9}; Peripheral membrane protein {ECO:0000250 | UniProtKB:Q8VEJ9}. Midbody Cytoplasm, cytoskeleton, spindle Note=Membrane-associated in the prevacuolar endosomal compartment Localizes to the midbody of dividing cells, interaction with ZFYVE19/ANCHR mediates retention at midbody (PubMed:24814515) Localized in two distinct rings on either side of the Flemming body


Tissue Location

Ubiquitously expressed.


Images



WB Suggested Anti-VPS4A Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:312500
Positive Control: Human Muscle



Host: Rabbit
Target Name: VPS4A
Sample Tissue: Human Fetal Heart
Antibody Dilution: 1.0µg/ml



Host: Rabbit
Target Name: VPS4A
Sample Tissue: Human Fetal Lung
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.