

FAM98A antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI13766

Product Information

Application WB
Primary Accession Q8NCA5

Other Accession <u>NM 015475</u>, <u>NP 056290</u>

Reactivity Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine

Predicted Human, Mouse, Rabbit, Chicken, Dog, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 55273

Additional Information

Gene ID 25940

Alias Symbol DKFZP564F0522, DKFZp686O03192

Other Names Protein FAM98A, FAM98A

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-FAM98A antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions FAM98A antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name FAM98A (<u>HGNC:24520</u>)

Function Positively stimulates PRMT1-induced protein arginine methylation

(PubMed:<u>28040436</u>). Involved in skeletal homeostasis (By similarity). Positively regulates lysosome peripheral distribution and ruffled border

formation in osteoclasts (By similarity).

Tissue Location Expressed strongly in colorectal cancer cells (PubMed:28040436). Expressed


strongly in colorectal cancer tissues compared to wild-type colon samples (at protein level) (PubMed:28040436). Expressed strongly in colorectal cancer

tissues compared to wild-type colon samples (PubMed:28040436)

References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

Images



WB Suggested Anti-FAM98A Antibody Titration: 0.2-1 $\mu g/ml$ Positive Control: COLO205 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.