

NDOR1 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI13894

Product Information

Application	WB
Primary Accession	Q9UHB4
Other Accession	XM_001716570 , XP_001716622
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66763

Additional Information

Gene ID	27158
Alias Symbol	MGC138148, NR1, bA350O14.9
Other Names	NADPH-dependent diflavin oxidoreductase 1 {ECO:0000255 HAMAP-Rule:MF_03178}, 1.6.-.- {ECO:0000255 HAMAP-Rule:MF_03178}, NADPH-dependent FMN and FAD-containing oxidoreductase {ECO:0000255 HAMAP-Rule:MF_03178}, Novel reductase 1, NDOR1 {ECO:0000255 HAMAP-Rule:MF_03178}, NR1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-NDOR1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	NDOR1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NDOR1 {ECO:0000255 HAMAP-Rule:MF_03178}
Function	NADPH-dependent reductase which is a central component of the cytosolic iron-sulfur (Fe-S) protein assembly (CIA) machinery (PubMed: 10625700 , PubMed: 15900210 , PubMed: 20802492 , PubMed: 23596212 , PubMed: 28648056). Transfers electrons from NADPH via its FAD and FMN prosthetic groups to the [2Fe-2S] cluster of CIAPIN1, another key component of the CIA machinery (PubMed: 20802492 , PubMed: 23596212 , PubMed: 28648056). In turn, this reduced cluster provides electrons for assembly of cytosolic iron-sulfur cluster proteins (PubMed: 20802492 ,

PubMed:[23596212](#)). It can also reduce the [2Fe-2S] cluster of CISD1 and activate this protein implicated in Fe/S cluster repair (PubMed:[28648056](#)). In vitro can fully activate methionine synthase/MTR in the presence of soluble cytochrome b5/CYB5A (PubMed:[12871938](#)).


Cellular Location

Cytoplasm, perinuclear region {ECO:0000255 | HAMAP- Rule:MF_03178, ECO:0000269 | PubMed:10625700, ECO:0000269 | PubMed:12871939}.
Note=Concentrated in perinuclear structure.
{ECO:0000255 | HAMAP-Rule:MF_03178, ECO:0000269 | PubMed:12871939}

Tissue Location

Low expression in brain, heart, kidney, pancreas, prostate and skeletal muscle. Highest levels in the placenta. Expressed in cancer cell lines including promyelocytic leukemia, HeLaS3, chronic myelogenous leukemia, lymphoblastic leukemia, Burkitt's lymphoma, colorectal adenocarcinoma, lung carcinoma, and melanoma G-361

Images



WB Suggested Anti-NDOR1 Antibody Titration: 1.0 µg/ml
Positive Control: ACHN Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.