

PARVG Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14157

Product Information

Application	WB
Primary Accession	Q9HBI0
Other Accession	XP_005261759
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37485

Additional Information

Gene ID	64098
Alias Symbol Other Names	PARVG, Gamma-parvin, PARVG
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 µl of distilled water. Final Anti-PARVG antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	PARVG Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	PARVG
Function	Plays a role with ILK in promoting the cell adhesion and spreading of leukocytes.
Cellular Location	Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side Cytoplasm, cytoskeleton. Note=Constituent of focal adhesions.
Tissue Location	Expressed predominantly in lymphoid organs, including spleen, thymus, lymph node, bone marrow and peripheral blood leukocytes and moderately in the digestive tract, including stomach, duodenum, jejunum, ileum, ileocecum and appendix, as well as in lung and liver. Also expressed in

tumors, but at a lower level than in the corresponding normal tissues.

References

Olski T.M.,et al.J. Cell Sci. 114:525-538(2001).
Collins J.E.,et al.Genome Res. 13:27-36(2003).
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham I.,et al.Nature 402:489-495(1999).

Images



Host: Rabbit
Target Name: PARVG
Sample Tissue: Large intestine Tumor lysates
Antibody Dilution: 1µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.