

# VPS16 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14184

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q9H269</a>
<b>Other Accession</b>	<a href="#">NM_080414</a> , <a href="#">NP_536339</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
<b>Predicted</b>	Human, Mouse, Rat, Rabbit, Dog, Horse, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	94694

## Additional Information

---

<b>Gene ID</b>	64601
<b>Alias Symbol</b>	hVPS16
<b>Other Names</b>	Vacuolar protein sorting-associated protein 16 homolog, hVPS16, VPS16
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-VPS16 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	VPS16 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	VPS16
<b>Function</b>	Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Believed to act as a core component of the putative HOPS and CORVET endosomal tethering complexes which are proposed to be involved in the Rab5-to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes. The CORVET complex is proposed to function as a Rab5 effector to mediate early endosome fusion probably in specific endosome subpopulations

(PubMed:[11382755](#), PubMed:[23351085](#), PubMed:[24554770](#), PubMed:[25266290](#), PubMed:[25783203](#)). Required for recruitment of VPS33A to the HOPS complex (PubMed:[23901104](#)). Required for fusion of endosomes and autophagosomes with lysosomes; the function is dependent on its association with VPS33A but not VPS33B (PubMed:[25783203](#)). The function in autophagosome- lysosome fusion implicates STX17 but not UVRAG (PubMed:[24554770](#)).

### Cellular Location


Late endosome membrane; Peripheral membrane protein; Cytoplasmic side. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome. Cytoplasmic vesicle, clathrin-coated vesicle {ECO:0000250|UniProtKB:Q920Q4}. Cytoplasmic vesicle, autophagosome. Note=Colocalizes with AP- 3, clathrin, Rab5 and Rab7b (By similarity). Cytoplasmic, peripheral membrane protein associated with early endosomes and late endosomes/lysosomes. {ECO:0000250|UniProtKB:Q920Q4, ECO:0000305}

### Tissue Location

Ubiquitous.

## Images

---



WB Suggested Anti-VPS16 Antibody Titration: 0.2-1 µg/ml  
Positive Control: MCF7 cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.