10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BCAM antibody - C-terminal region

Rabbit Polyclonal Antibody
Catalog # AI14503

Product Information

Application	WB
Primary Accession	P50895
Other Accession	NM_005581, NP_005572
Reactivity	Human, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Rabbit, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67405

Additional Information

Gene ID	4059
Alias Symbol	AU, CD239, LU, MSK19
Other Names	Basal cell adhesion molecule, Auberger B antigen, B-CAM cell surface glycoprotein, F8/G253 antigen, Lutheran antigen, Lutheran blood group glycoprotein, CD239, BCAM, LU, MSK19
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-BCAM antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	BCAM antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	BCAM
Synonyms	LU, MSK19
Function	Transmembrane glycoprotein that functions as both a receptor and an adhesion molecule playing a crucial role in cell adhesion, motility, migration and invasion (PubMed:9616226, PubMed:31413112). Extracellular domain enables binding to extracellular matrix proteins, such as laminin, integrin and other ligands while its intracellular domain interacts with cytoskeletal proteins like hemoglobin, facilitating cell signal transduction (PubMed:17158232). Serves as a receptor for laminin alpha-5/LAMA5 to promote cell adhesion

	(PubMed:15975931). Mechanistically, JAK2 induces BCAM phosphorylation and activates its adhesion to laminin by stimulating a Rap1/AKT signaling pathway in the absence of EPOR (PubMed:23160466).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Wide tissue distribution (highest in the pancreas and very low in brain). Closely associated with the basal layer of cells in epithelia and the endothelium of blood vessel walls

References

Parsons S.F.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:5496-5500(1995).
Grimwood J.,et al.Nature 428:529-535(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Campbell I.G.,et al.Cancer Res. 54:5761-5765(1994).
Wang C.,et al.Submitted (NOV-2007) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-BCAM Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:1562500
Positive Control: HepG2 cell lysate
BCAM is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.