

ACSM3 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI14570

Product Information

В
5 <u>3FZ2</u>
<u>M_005622, NP_005613</u>
ıman, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
ıman, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Guinea Pig, Horse, Bovine
bbit
lyclonal
153

Additional Information


Gene ID	6296
Alias Symbol Other Names	SA, SAH Acyl-coenzyme A synthetase ACSM3, mitochondrial, 6.2.1.2, Acyl-CoA synthetase medium-chain family member 3, ButyrateCoA ligase 3, Butyryl-coenzyme A synthetase 3, Middle-chain acyl-CoA synthetase 3, Protein SA homolog, ACSM3, SAH
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-ACSM3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	ACSM3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACSM3
Synonyms	SAH
Function	Catalyzes the activation of fatty acids by CoA to produce an acyl-CoA, the first step in fatty acid metabolism (PubMed: <u>11772874</u>). Capable of activating medium-chain fatty acids with a preference for isobutyrate among fatty acids with 2-6 carbon atoms (By similarity).
Cellular Location	Mitochondrion. Mitochondrion matrix {ECO:0000250 UniProtKB:Q3UNX5}

References

Iwai N., et al. Hypertension 23:375-380(1994). Nabika T., et al. Hypertension 25:6-13(1995). Loftus B.J., et al. Genomics 60:295-308(1999). Suzuki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases. Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.