

AKR1D1 antibody - middle region

Rabbit Polyclonal Antibody Catalog # AI14572

Product Information

Application WB Primary Accession P51857

Other Accession NM 005989, NP 005980

Reactivity Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine

Predicted Human, Mouse, Rat, Chicken, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 37377

Additional Information

Gene ID 6718

Alias Symbol 3o5bred, CBAS2, SRD5B1

Other Names 3-oxo-5-beta-steroid 4-dehydrogenase, 1.3.1.3, Aldo-keto reductase family 1

member D1, Delta(4)-3-ketosteroid 5-beta-reductase, Delta(4)-3-oxosteroid

5-beta-reductase, AKR1D1, SRD5B1

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-AKR1D1 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions AKR1D1 antibody - middle region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name AKR1D1

Synonyms SRD5B1

Function Catalyzes the stereospecific NADPH-dependent reduction of the C4-C5

double bond of bile acid intermediates and steroid hormones carrying a delta(4)-3-one structure to yield an A/B cis-ring junction. This cis-configuration is crucial for bile acid biosynthesis and plays important roles in steroid metabolism. Capable of reducing a broad range of delta-(4)-3-ketosteroids from C18 (such as, 17beta- hydroxyestr-4-en-3-one) to C27 (such as,

7alpha-hydroxycholest-4-en-3- one).


Cellular Location Cytoplasm.

Tissue Location Highly expressed in liver. Expressed in testis and weakly in colon.

References

Kondo K.-H.,et al.Eur. J. Biochem. 219:357-363(1994). Charbonneau A.,et al.Biochim. Biophys. Acta 1517:228-235(2001). Ota T.,et al.Nat. Genet. 36:40-45(2004). Hillier L.W.,et al.Nature 424:157-164(2003). Scherer S.W.,et al.Science 300:767-772(2003).

Images



WB Suggested Anti-AKR1D1 Antibody Titration: 1.0 µg/ml Positive Control: RPMI-8226 Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.