

STX8 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI14614

Product Information

Application WB
Primary Accession Q9UNK0

Other Accession <u>NM 004853</u>, <u>NP 004844</u>

ReactivityHuman, Mouse, Rat, Zebrafish, Dog, Guinea Pig, Horse, Bovine **Predicted**Human, Mouse, Rat, Zebrafish, Chicken, Dog, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 26907

Additional Information

Gene ID 9482

Alias Symbol CARB

Other Names Syntaxin-8, STX8

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-STX8 antibody concentration is 1 mg/ml

in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C.

Avoid repeat freeze-thaw cycles.

Precautions STX8 antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name STX8

Function Vesicle trafficking protein that functions in the early secretory pathway,

possibly by mediating retrograde transport from cis- Golgi membranes to the

ER.

Cellular Location Membrane; Single-pass type IV membrane protein. Note=Preferentially

associated with the early endosome. To a lesser extent, also present in late

endosome, the plasma membrane and coated pits (By similarity).


Tissue Location Highly expressed in heart. Also found in brain, kidney, liver, lung, placenta,

skeletal muscle, spleen and pancreas

References

Steegmaier M., et al.J. Biol. Chem. 273:34171-34179(1998).
Thoreau V., et al. Biochem. Biophys. Res. Commun. 257:577-583(1999).
Subramaniam V.N., et al. Submitted (DEC-1997) to the EMBL/GenBank/DDBJ databases.
Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-STX8 Antibody Titration: 1.0 µg/ml Positive Control: Fetal Heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.