

# ACSS2 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14617

## Product Information

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">Q9NR19</a>
<b>Other Accession</b>	<a href="#">NM_001076552</a> , <a href="#">NP_001070020</a>
<b>Reactivity</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Goat, Dog, Guinea Pig, Horse, Bovine
<b>Predicted Host</b>	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
<b>Clonality</b>	Rabbit
<b>Calculated MW</b>	Polyclonal 78580

## Additional Information

<b>Gene ID</b>	55902
<b>Alias Symbol</b> <b>Other Names</b>	ACAS2, ACECS, ACS, ACSA, DKFZp762G026, dj1161H23.1 Acetyl-coenzyme A synthetase, cytoplasmic, 6.2.1.1, Acetate--CoA ligase, Acetyl-CoA synthetase, ACS, AceCS, Acyl-CoA synthetase short-chain family member 2, Acyl-activating enzyme, ACSS2, ACAS2
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-ACSS2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	ACSS2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

<b>Name</b>	ACSS2
<b>Synonyms</b>	ACAS2
<b>Function</b>	Catalyzes the synthesis of acetyl-CoA from short-chain fatty acids (PubMed: <a href="#">10843999</a> , PubMed: <a href="#">28003429</a> , PubMed: <a href="#">28552616</a> ). Acetate is the preferred substrate (PubMed: <a href="#">10843999</a> , PubMed: <a href="#">28003429</a> ). Can also utilize propionate with a much lower affinity (By similarity). Nuclear ACSS2 promotes glucose deprivation-induced lysosomal biogenesis and autophagy, tumor cell survival and brain tumorigenesis (PubMed: <a href="#">28552616</a> ). Glucose deprivation

results in AMPK-mediated phosphorylation of ACSS2 leading to its translocation to the nucleus where it binds to TFEB and locally produces acetyl-CoA for histone acetylation in the promoter regions of TFEB target genes thereby activating their transcription (PubMed:[28552616](#)). The regulation of genes associated with autophagy and lysosomal activity through ACSS2 is important for brain tumorigenesis and tumor survival (PubMed:[28552616](#)). Acts as a chromatin-bound transcriptional coactivator that up-regulates histone acetylation and expression of neuronal genes (By similarity). Can be recruited to the loci of memory-related neuronal genes to maintain a local acetyl-CoA pool, providing the substrate for histone acetylation and promoting the expression of specific genes, which is essential for maintaining long-term spatial memory (By similarity).

### Cellular Location

Cytoplasm, cytosol. Cytoplasm {ECO:0000250|UniProtKB:Q9QXG4}. Nucleus Note=Glucose deprivation results in its AMPK-dependent phosphorylation and subsequent nuclear translocation (PubMed:28552616). Phosphorylation at Ser-659, leads to exposure of its nuclear localization signal which is required for its interaction with KPNA1 and subsequent translocation to the nucleus (PubMed:28552616). Found in the cytoplasm in undifferentiated neurons and upon differentiation, translocates to nucleus (By similarity). {ECO:0000250|UniProtKB:Q9QXG4, ECO:0000269|PubMed:28552616}

### References

Luong A.,et al.J. Biol. Chem. 275:26458-26466(2000).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Deloukas P.,et al.Nature 414:865-871(2001).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Zahedi R.P.,et al.J. Proteome Res. 7:526-534(2008).

### Images



WB Suggested Anti-ACSS2 Antibody Titration: 1.0 µg/ml  
Positive Control: Fetal Heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.