10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PRTN3 antibody - N-terminal region

Rabbit Polyclonal Antibody
Catalog # AI14624

Product Information

Application	WB
Primary Accession	P24158
Other Accession	NM_002777, NP_002768
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27807

Additional Information

Gene ID	5657
Alias Symbol	ACPA, AGP7, C-ANCA, CANCA, MBN, MBT, NP4, P29, PR-3, PR3
Other Names	Myeloblastin, 3.4.21.76, AGP7, C-ANCA antigen, Leukocyte proteinase 3, PR-3, PR3, Neutrophil proteinase 4, NP-4, P29, Wegener autoantigen, PRTN3, MBN
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-PRTN3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	PRTN3 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PRTN3
Synonyms	MBN
Function	Serine protease that degrades elastin, fibronectin, laminin, vitronectin, and collagen types I, III, and IV (in vitro) (PubMed:2033050, PubMed:28240246, PubMed:3198760). By cleaving and activating receptor F2RL1/PAR-2, enhances endothelial cell barrier function and thus vascular integrity during neutrophil transendothelial migration (PubMed:23202369). Plays a role in neutrophil transendothelial migration, probably when associated with CD177 (PubMed:22266279). Triggers inflammatory processes in neutrophils by interacting with ADGRG3 upstream of F2RL1/PAR2 activation

(PubMed:36302784).

Cellular Location

Cytoplasmic granule. Secreted. Cell membrane; Peripheral membrane protein; Extracellular side. Membrane raft; Peripheral membrane protein; Extracellular side. Note=Localizes predominantly to azurophil granules (primary secretory granules) in neutrophils (PubMed:18462208, PubMed:2033050, PubMed:3198760, PubMed:7897245) Secreted upon neutrophil stimulation by TNF, lipopolysaccharide (LPS), fMLP and CXCL8/IL8 or during neutrophil transmigration (PubMed:22266279, PubMed:28240246). Following secretion tethered to the cell membrane by CD177 (PubMed:18462208, PubMed:22266279)


Tissue Location

Expressed in polymorphonuclear leukocytes (at protein level) (PubMed:2033050, PubMed:3198760, PubMed:7897245) Expressed in neutrophils (at protein level) (PubMed:17244676, PubMed:18462208, PubMed:21193407, PubMed:22266279, PubMed:28240246) Expressed in differentiating neutrophils (PubMed:18462208)

References

Labbaye C.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:9253-9256(1991).
Grimwood J.,et al.Nature 428:529-535(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Zimmer M.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:8215-8219(1992).
Sturrock A.B.,et al.J. Biol. Chem. 267:21193-21199(1992).

Images



WB Suggested Anti-PRTN3 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Lung

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.