

PTGDR antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14652

Product Information

Application	WB
Primary Accession	Q13258
Other Accession	NM_000953 , NP_000944
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40271

Additional Information

Gene ID	5729
Alias Symbol Other Names	AS1, ASRT1, DP, DP1, MGC49004, PTGDR1 Prostaglandin D2 receptor, PGD receptor, PGD2 receptor, Prostanoid DP receptor, PTGDR
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-PTGDR antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	PTGDR antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PTGDR
Function	Receptor for prostaglandin D2 (PGD2). The activity of this receptor is mainly mediated by G(s) proteins that stimulate adenylate cyclase, resulting in an elevation of intracellular cAMP. A mobilization of calcium is also observed, but without formation of inositol 1,4,5-trisphosphate (By similarity). Involved in PLA2G3- dependent maturation of mast cells. PLA2G3 is secreted by immature mast cells and acts on nearby fibroblasts upstream to PTDGS to synthesize PGD2, which in turn promotes mast cell maturation and degranulation via PTGDR (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein

Tissue Location

Expressed in retinal choroid, ciliary epithelium, longitudinal and circular ciliary muscles, iris, small intestine and platelet membranes.

References

Boie Y.,et al.J. Biol. Chem. 270:18910-18916(1995).


Martin A.L.,et al.Submitted (APR-2007) to the EMBL/GenBank/DDBJ databases.

Heilig R.,et al.Nature 421:601-607(2003).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Town M.H.,et al.Prostaglandins 25:13-28(1983).

Images



WB Suggested Anti-PTGDR Antibody Titration: 1.0 µg/ml
Positive Control: THP-1 Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.