

# DTNA antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14676

## Product Information


|                          |                                                             |
|--------------------------|-------------------------------------------------------------|
| <b>Application</b>       | WB                                                          |
| <b>Primary Accession</b> | <a href="#">B4DIRO</a>                                      |
| <b>Other Accession</b>   | <a href="#">NM_001198942</a> , <a href="#">NP_001185871</a> |
| <b>Reactivity</b>        | Human, Mouse, Rat, Dog, Guinea Pig, Horse, Bovine, Yeast    |
| <b>Predicted</b>         | Human, Mouse, Rat, Chicken, Dog, Guinea Pig, Horse, Bovine  |
| <b>Host</b>              | Rabbit                                                      |
| <b>Clonality</b>         | Polyclonal                                                  |
| <b>Calculated MW</b>     | 49 kDa                                                      |

## Additional Information

|                                     |                                                                                                                                                                                               |
|-------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Alias Symbol</b>                 | D18S892E, DRP3, DTN, FLJ96209, LVNC1                                                                                                                                                          |
| <b>Format</b>                       | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.                                                                                             |
| <b>Reconstitution &amp; Storage</b> | Add 50 ul of distilled water. Final anti-DTNA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles. |
| <b>Precautions</b>                  | DTNA antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.                                                                           |

## Protein Information

## Images



WB Suggested Anti-DTNA Antibody Titration: 1.0 µg/ml  
Positive Control: Hela Whole Cell