

# CUL4A antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI14702

## Product Information

Application	WB
Primary Accession	<a href="#">Q13619</a>
Other Accession	<a href="#">NM_001008895</a> , <a href="#">NP_001008895</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87680

## Additional Information

Gene ID	8451
Other Names	Cullin-4A, CUL-4A, CUL4A
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-CUL4A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	CUL4A antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	CUL4A {ECO:0000303   PubMed:9721878, ECO:0000312   HGNC:HGNC:2554}
Function	Core component of multiple cullin-RING-based E3 ubiquitin- protein ligase complexes which mediate the ubiquitination of target proteins (PubMed: <a href="#">14578910</a> , PubMed: <a href="#">14739464</a> , PubMed: <a href="#">15448697</a> , PubMed: <a href="#">15548678</a> , PubMed: <a href="#">15811626</a> , PubMed: <a href="#">16678110</a> , PubMed: <a href="#">17041588</a> , PubMed: <a href="#">24209620</a> , PubMed: <a href="#">30166453</a> , PubMed: <a href="#">33854232</a> , PubMed: <a href="#">33854239</a> ). As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed: <a href="#">14578910</a> , PubMed: <a href="#">14739464</a> , PubMed: <a href="#">15448697</a> , PubMed: <a href="#">15548678</a> , PubMed: <a href="#">15811626</a> , PubMed: <a href="#">16678110</a> , PubMed: <a href="#">17041588</a> , PubMed: <a href="#">24209620</a> ). The E3 ubiquitin- protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with

TIP120A/CAND1 (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). DCX(DET1-COP1) directs ubiquitination of JUN (PubMed:[14739464](#)). DCX(DDB2) directs ubiquitination of XPC (PubMed:[15811626](#)). DCX(DDB2) ubiquitinates histones H3-H4 and is required for efficient histone deposition during replication-coupled (H3.1) and replication-independent (H3.3) nucleosome assembly, probably by facilitating the transfer of H3 from ASF1A/ASF1B to other chaperones involved in histone deposition (PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of p53/TP53 in response to radiation-induced DNA damage and during DNA replication (PubMed:[14578910](#), PubMed:[15448697](#), PubMed:[15548678](#)). DCX(DTL) directs autoubiquitination of DTL (PubMed:[23478445](#)). In association with DDB1 and SKP2 probably is involved in ubiquitination of CDKN1B/p27kip (PubMed:[16537899](#)). Is involved in ubiquitination of HOXA9 (PubMed:[14609952](#)). The DDB1-CUL4A- DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (PubMed:[26431207](#)). The DCX(ERCC8) complex (also named CSA complex) plays a role in transcription-coupled repair (TCR) (PubMed:[12732143](#), PubMed:[32355176](#), PubMed:[38316879](#)). A number of DCX complexes (containing either TRPC4AP or DCAF12 as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:[29779948](#)). The DCX(AMBRA1) complex is a master regulator of the transition from G1 to S cell phase by mediating ubiquitination of phosphorylated cyclin-D (CCND1, CCND2 and CCND3) (PubMed:[33854232](#), PubMed:[33854239](#)). The DCX(AMBRA1) complex also acts as a regulator of Cul5-RING (CRL5) E3 ubiquitin-protein ligase complexes by mediating ubiquitination and degradation of Elongin-C (ELOC) component of CRL5 complexes (PubMed:[30166453](#)). With CUL4B, contributes to ribosome biogenesis (PubMed:[26711351](#)).

## References


---

Chen L.-C.,et al.Cancer Res. 58:3677-3683(1998).  
Higa L.A.,et al.Nat. Cell Biol. 5:1008-1015(2003).  
Matsuda N.,et al.DNA Repair 4:537-545(2005).  
Dunham A.,et al.Nature 428:522-528(2004).  
Osaka F.,et al.Genes Dev. 12:2263-2268(1998).


## Images

---

WB Suggested Anti-CUL4A Antibody Titration: 1.0 µg/ml  
Positive Control: Fetal Heart



### CUL4A



Rabbit Anti-CUL4A  
Sample Type: Human Fetal Lung  
Antibody Concentration: 1ug/mL

Host: Rabbit  
Target Name: CUL4A  
Sample Tissue: Human Fetal Lung  
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.