

# RPS9 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI14718

#### **Product Information**

Application WB Primary Accession P46781

Other Accession NM 001013, NP 001004

**Reactivity** Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine

**Predicted** Human, Rabbit, Zebrafish, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 22591

#### **Additional Information**

**Gene ID** 6203

Alias Symbol S9

Other Names 40S ribosomal protein S9, RPS9

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-RPS9 antibody concentration is 1 mg/ml

in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C.

Avoid repeat freeze-thaw cycles.

**Precautions** RPS9 antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name RPS9 ( HGNC:10442)

**Function** Component of the small ribosomal subunit (PubMed: <u>23636399</u>). The

ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed: 23636399). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with

the nascent pre-rRNA and work in concert to generate RNA folding,

modifications, rearrangements and cleavage as well as targeted degradation


of pre-ribosomal RNA by the RNA exosome (PubMed:34516797).

**Cellular Location** Cytoplasm. Nucleus, nucleolus. Note=Localized in cytoplasmic mRNP granules

### References

Frigerio J.-M.,et al.Biochim. Biophys. Acta 1262:64-68(1995). Yoshihama M.,et al.Genome Res. 12:379-390(2002). Grimwood J.,et al.Nature 428:529-535(2004). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Vladimirov S.N.,et al.Eur. J. Biochem. 239:144-149(1996).

## **Images**



WB Suggested Anti-RPS9 Antibody Titration: 1.0  $\mu$ g/ml Positive Control: HepG2 Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.