

CYP2U1 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI14742

Product Information

Application	WB
Primary Accession	Q7Z449
Other Accession	NM_183075 , NP_898898
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Goat, Dog, Horse, Bovine, Sheep
Predicted	Human, Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61987

Additional Information

Gene ID	113612
Alias Symbol	P450TEC
Other Names	Cytochrome P450 2U1, 1.14.14.1, CYP2U1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-CYP2U1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	CYP2U1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CYP2U1 {ECO:0000303 PubMed:14660610, ECO:0000312 HGNC:HGNC:20582}
Function	A cytochrome P450 monooxygenase involved in the metabolism of arachidonic acid and its conjugates (PubMed: 14660610 , PubMed: 24563460). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed: 14660610 , PubMed: 24563460). Acts as an omega and omega-1 hydroxylase for arachidonic acid and possibly for other long chain fatty acids. May modulate the arachidonic acid signaling pathway and play a role in other fatty acid signaling processes (PubMed: 14660610 , PubMed: 24563460). May down-regulate the biological

activities of N-arachidonoyl-serotonin, an endocannabinoid that has anti-nociceptive effects through inhibition of fatty acid amide hydrolase FAAH, TRPV1 receptor and T-type calcium channels. Catalyzes C-2 oxidation of the indole ring of N-arachidonoyl-serotonin forming a less active product 2-oxo-N-arachidonoyl-serotonin (PubMed:[24563460](#)).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein. Microsome membrane; Multi- pass membrane protein. Mitochondrion inner membrane; Multi-pass membrane protein


Tissue Location

Widely expressed with stronger expression in thymus, heart and cerebellum.

References

Chuang S.S.,et al.J. Biol. Chem. 279:6305-6314(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Karlgren M.,et al.Biochem. Biophys. Res. Commun. 315:679-685(2004).
Choudhary D.,et al.Arch. Biochem. Biophys. 436:50-61(2005).
Tesson C.,et al.Am. J. Hum. Genet. 91:1051-1064(2012).

Images



WB Suggested Anti-CYP2U1 Antibody Titration: 1.0 µg/ml
Positive Control: Jurkat Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.