10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PPP2R4 antibody - C-terminal region

Rabbit Polyclonal Antibody
Catalog # AI14766

Product Information

Application	WB
Primary Accession	Q15257
Other Accession	NM_178001, NP_821068
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Chicken, Dog, Guinea Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40668

Additional Information

Gene ID	5524
Alias Symbol	MGC2184, PP2A, PR53, PTPA
Other Names	Serine/threonine-protein phosphatase 2A activator, 5.2.1.8, PP2A, subunit B', PR53 isoform, Phosphotyrosyl phosphatase activator, PTPA, Serine/threonine-protein phosphatase 2A regulatory subunit 4, Serine/threonine-protein phosphatase 2A regulatory subunit B', PPP2R4, PTPA
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-PPP2R4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	PPP2R4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PTPA (HGNC:9308)
Synonyms	PPP2R4
Function	PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides (By similarity). Acts as a regulatory subunit for serine/threonine-protein phosphatase 2A (PP2A) (PubMed:16916641, PubMed:36073231). Modulates PP2A activity or substrate specificity, probably by inducing a conformational change in the catalytic subunit, a proposed direct target of the PPIase (PubMed:16916641).

Can reactivate inactive phosphatase PP2A-phosphatase methylesterase complexes (PP2A(i)) in presence of ATP and Mg(2+) (By similarity). Reversibly stimulates the variable phosphotyrosyl phosphatase activity of PP2A core heterodimer PP2A(D) in presence of ATP and Mg(2+) (in vitro) (PubMed:16916641). The phosphotyrosyl phosphatase activity is dependent of an ATPase activity of the PP2A(D):PPP2R4 complex (PubMed:16916641). Is involved in apoptosis; the function appears to be independent from PP2A (PubMed:17333320).


Cellular Location Cytoplasm. Nucleus

Tissue Location Widely expressed.

References

Cayla X.,et al.J. Biol. Chem. 269:15668-15675(1994).
Van Hoof C.,et al.Genomics 28:261-272(1995).
Janssens V.,et al.Eur. J. Biochem. 267:4406-4413(2000).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-PPP2R4 Antibody Titration: 1.0 µg/ml
Positive Control: 721_B Whole CellPPP2R4 is strongly supported by BioGPS gene expression data to be expressed in Human 721_B cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.