

# VPS33A Antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI14769

## Product Information

|                          |                                                                           |
|--------------------------|---------------------------------------------------------------------------|
| <b>Application</b>       | WB                                                                        |
| <b>Primary Accession</b> | <a href="#">Q96AX1</a>                                                    |
| <b>Other Accession</b>   | <a href="#">NM_022916</a> , <a href="#">NP_075067</a>                     |
| <b>Reactivity</b>        | Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine |
| <b>Predicted</b>         | Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine |
| <b>Host</b>              | Rabbit                                                                    |
| <b>Clonality</b>         | Polyclonal                                                                |
| <b>Calculated MW</b>     | 67611                                                                     |

## Additional Information

|                                     |                                                                                                                                                                                                 |
|-------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>                      | 65082                                                                                                                                                                                           |
| <b>Other Names</b>                  | Vacuolar protein sorting-associated protein 33A, hVPS33A, VPS33A                                                                                                                                |
| <b>Format</b>                       | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.                                                                                               |
| <b>Reconstitution &amp; Storage</b> | Add 50 ul of distilled water. Final anti-VPS33A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles. |
| <b>Precautions</b>                  | VPS33A Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.                                                                               |

## Protein Information

|                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Name</b>     | VPS33A                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Function</b> | Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Believed to act as a core component of the putative HOPS and CORVET endosomal tethering complexes which are proposed to be involved in the Rab5-to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic towards lysosomes. The CORVET complex is proposed to function as a Rab5 effector to mediate early endosome fusion probably in specific endosome subpopulations (PubMed: <a href="#">23351085</a> , PubMed: <a href="#">24554770</a> , PubMed: <a href="#">25266290</a> , |

PubMed:[25783203](#)). Required for fusion of endosomes and autophagosomes with lysosomes; the function is dependent on its association with VPS16 but not VIPAS39 (PubMed:[25783203](#)). The function in autophagosome-lysosome fusion implicates STX17 but not UVRAG (PubMed:[24554770](#)).

## Cellular Location

Cytoplasmic vesicle {ECO:0000250 | UniProtKB:Q63615}. Late endosome membrane; Peripheral membrane protein; Cytoplasmic side. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome. Cytoplasmic vesicle, autophagosome. Cytoplasmic vesicle, clathrin-coated vesicle


## References

---

Suzuki T.,et al.Proc. Natl. Acad. Sci. U.S.A. 100:1146-1150(2003).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Zhu G.D.,et al.Mol. Biol. Cell 20:1223-1240(2009).  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

## Images

---



Host: Rabbit  
Target Name: VPS33A  
Sample Tissue: 721\_B whole cell lysate  
S  
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.