10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PPP6C Antibody - C-terminal region

Rabbit Polyclonal Antibody
Catalog # AI14822

Product Information

Application	WB
Primary Accession	O00743
Other Accession	NP_002712
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35144

Additional Information

Gene ID	5537
Alias Symbol	PPP6C, PPP6,
Other Names	Serine/threonine-protein phosphatase 6 catalytic subunit, PP6C, 3.1.3.16, Serine/threonine-protein phosphatase 6 catalytic subunit, N-terminally processed, PPP6C, PPP6
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 &mu, l of distilled water. Final Anti-PPP6C antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	PPP6C Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPP6C {ECO:0000303|PubMed:29053956, ECO:0000312|HGNC:HGNC:9323}
Function	Catalytic subunit of protein phosphatase 6 (PP6) (PubMed: 17079228, PubMed:29053956, PubMed:32474700). PP6 is a component of a signaling pathway regulating cell cycle progression in response to IL2 receptor stimulation (PubMed:10227379). N-terminal domain restricts G1 to S phase progression in cancer cells, in part through control of cyclin D1 (PubMed:17568194). During mitosis, regulates spindle positioning (PubMed:27335426). Down-regulates MAP3K7 kinase activation of the IL1 signaling pathway by dephosphorylation of MAP3K7 (PubMed:17079228). Also participates in the innate immune defense against viruses by desphosphorylating RIGI, an essential step that triggers RIGI-mediated

	signaling activation (PubMed:29053956). Also regulates innate immunity by acting as a negative regulator of the cGAS-STING pathway: mediates dephosphorylation and inactivation of CGAS and STING1 (PubMed:32474700, PubMed:32753499). CGAS dephosphorylation at 'Ser-435' impairs its ability to bind GTP, thereby inactivating it (PubMed:32474700).
Cellular Location	Mitochondrion. Cytoplasm
Tissue Location	Ubiquitously expressed in all tissues tested with highest expression levels in testis, heart, kidney, brain, stomach, liver and skeletal muscle and lowest in placenta, lung colon and spleen.

References

Bastians H.,et al.J. Cell Sci. 109:2865-2874(1996).
Filali M.,et al.J. Cell. Biochem. 73:153-163(1999).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Humphray S.J.,et al.Nature 429:369-374(2004).

Images



Host: Rabbit
Target Name: PPP6C
Sample Tissue: Hela Whole Cell lysates
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.