

HIP1 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI14988

Product Information

Application	WB
Primary Accession	O00291
Other Accession	NP_005329
Reactivity	Human
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116221

Additional Information

Gene ID	3092
Alias Symbol Other Names	HIP1, Huntingtin-interacting protein 1, HIP-1, Huntingtin-interacting protein I, HIP-I, HIP1
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 µl of distilled water. Final Anti-HIP1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	HIP1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HIP1
Function	Plays a role in clathrin-mediated endocytosis and trafficking (PubMed: 11532990 , PubMed: 11577110 , PubMed: 11889126). Involved in regulating AMPA receptor trafficking in the central nervous system in an NMDA-dependent manner (By similarity). Regulates presynaptic nerve terminal activity (By similarity). Enhances androgen receptor (AR)- mediated transcription (PubMed: 16027218). May act as a proapoptotic protein that induces cell death by acting through the intrinsic apoptosis pathway (PubMed: 11007801). Binds 3-phosphoinositides (via ENTH domain) (PubMed: 14732715). May act through the ENTH domain to promote cell survival by stabilizing receptor tyrosine kinases following ligand-induced

endocytosis (PubMed:[14732715](#)). May play a functional role in the cell filament networks (PubMed:[18790740](#)). May be required for differentiation, proliferation, and/or survival of somatic and germline progenitors (PubMed:[11007801](#), PubMed:[12163454](#)).

Cellular Location

Cytoplasm. Nucleus. Endomembrane system. Cytoplasmic vesicle, clathrin-coated vesicle membrane. Note=Shuttles between cytoplasm and nucleus. Nuclear translocation can be induced by AR

Tissue Location

Ubiquitously expressed with the highest level in brain. Expression is up-regulated in prostate and colon cancer

References

Kim R.N.,et al.Submitted (JUL-2013) to the EMBL/GenBank/DDBJ databases.


Ota T.,et al.Nat. Genet. 36:40-45(2004).

Hillier L.W.,et al.Nature 424:157-164(2003).

Gervais F.G.,et al.Nat. Cell Biol. 4:95-105(2002).

Huq A.H.M.M.,et al.Submitted (MAR-1998) to the EMBL/GenBank/DDBJ databases.

Images



Host: Rabbit
Target Name: HIP1
Sample Tissue: A549 Whole Cell lysates
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.