

# IL3RA antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI15012

#### **Product Information**

Application WB Primary Accession P26951

Other Accession NM 002183, NP 002174

Reactivity Human
Predicted Human
Host Rabbit
Clonality Polyclonal
Calculated MW 43330

#### **Additional Information**

Gene ID 3563

Alias Symbol CD123, IL3R, IL3RAY, IL3RX, IL3RY, MGC34174, hIL-3Ra

Other Names Interleukin-3 receptor subunit alpha, IL-3 receptor subunit alpha, IL-3R

subunit alpha, IL-3R-alpha, IL-3RA, CD123, IL3RA, IL3R

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-IL3RA antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

**Precautions** IL3RA antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name IL3RA ( HGNC:6012)

Synonyms IL3R

**Function** Cell surface receptor for IL3 expressed on hematopoietic progenitor cells,

monocytes and B-lymphocytes that controls the production and

differentiation of hematopoietic progenitor cells into lineage-restricted cells (PubMed: 10527461). Ligand stimulation rapidly induces hetrodimerization with IL3RB, phosphorylation and enzyme activity of effector proteins such as

JAK2 and PI3K that play a role in signaling cell proliferation and


differentiation. Activation of JAK2 leads to STAT5-mediated transcriptional

program (By similarity).

### References

Kitamura T.,et al.Cell 66:1165-1174(1991). Kosugi H.,et al.Biochem. Biophys. Res. Commun. 208:360-367(1995). Chen J.,et al.J. Biol. Chem. 284:5763-5773(2009). Ota T.,et al.Nat. Genet. 36:40-45(2004). Ross M.T.,et al.Nature 434:325-337(2005).

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.