

PALB2 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15014

Product Information

Application	WB
Primary Accession	Q86YC2
Other Accession	NM_024675 , BAB15140
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	131295

Additional Information

Gene ID	79728
Alias Symbol	FANCN, PNCA3
Other Names	Partner and localizer of BRCA2, PALB2, FANCN
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-PALB2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	PALB2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PALB2
Synonyms	FANCN
Function	Plays a critical role in homologous recombination repair (HRR) through its ability to recruit BRCA2 and RAD51 to DNA breaks (PubMed: 16793542 , PubMed: 19369211 , PubMed: 19423707 , PubMed: 22941656 , PubMed: 24141787 , PubMed: 28319063). Strongly stimulates the DNA strand-invasion activity of RAD51, stabilizes the nucleoprotein filament against a disruptive BRC3-BRC4 polypeptide and helps RAD51 to overcome the suppressive effect of replication protein A (RPA) (PubMed: 20871615). Functionally cooperates with RAD51AP1 in promoting of D-loop formation by RAD51 (PubMed: 20871616). Serves as the molecular scaffold in the formation

of the BRCA1-PALB2-BRCA2 complex which is essential for homologous recombination (PubMed:[19369211](#)). Via its WD repeats is proposed to scaffold a HR complex containing RAD51C and BRCA2 which is thought to play a role in HR-mediated DNA repair (PubMed:[24141787](#)). Essential partner of BRCA2 that promotes the localization and stability of BRCA2 (PubMed:[16793542](#)). Also enables its recombinational repair and checkpoint functions of BRCA2 (PubMed:[16793542](#)). May act by promoting stable association of BRCA2 with nuclear structures, allowing BRCA2 to escape the effects of proteasome-mediated degradation (PubMed:[16793542](#)). Binds DNA with high affinity for D loop, which comprises single-stranded, double-stranded and branched DNA structures (PubMed:[20871616](#)). May play a role in the extension step after strand invasion at replication-dependent DNA double-strand breaks; together with BRCA2 is involved in both POLH localization at collapsed replication forks and DNA polymerization activity (PubMed:[24485656](#)).


Cellular Location

Nucleus Note=Colocalizes with BRCA2 and BRCA1 in nuclear foci

References

Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Martin J.,et al.Nature 432:988-994(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Xia B.,et al.Mol. Cell 22:719-729(2006).

Images



WB Suggested Anti-PALB2 Antibody Titration: 1.0 µg/ml
Positive Control: HeLa Whole Cell
PALB2 is supported by BioGPS gene expression data to be expressed in HeLa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.