

DYRK2 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI15090

Product Information

Application	WB
Primary Accession	<u>Q92630</u>
Other Accession	<u>NM_003583</u> , <u>NP_003574</u>
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66652
Host Clonality Calculated MW	Rabbit Polyclonal 66652

Additional Information

Gene ID	8445
Alias Symbol Other Names	FLJ21217, FLJ21365 Dual specificity tyrosine-phosphorylation-regulated kinase 2, 2.7.12.1, DYRK2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-DYRK2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	DYRK2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

NameDYRK2FunctionSerine/threonine-protein kinase involved in the regulation of the mitotic cell
cycle, cell proliferation, apoptosis, organization of the cytoskeleton and
neurite outgrowth. Functions in part via its role in ubiquitin-dependent
proteasomal protein degradation. Functions downstream of ATM and
phosphorylates p53/TP53 at 'Ser-46', and thereby contributes to the induction
of apoptosis in response to DNA damage. Phosphorylates NFATC1, and
thereby inhibits its accumulation in the nucleus and its transcription factor
activity. Phosphorylates EIF2B5 at 'Ser-544', enabling its subsequent
phosphorylation and inhibition by GSK3B. Likewise, phosphorylation of
NFATC1, CRMP2/DPYSL2 and CRMP4/DPYSL3 promotes their subsequent
phosphorylation by GSK3B. May play a general role in the priming of GSK3

	substrates. Inactivates GYS1 by phosphorylation at 'Ser-641', and potentially also a second phosphorylation site, thus regulating glycogen synthesis. Mediates EDVP E3 ligase complex formation and is required for the phosphorylation and subsequent degradation of KATNA1. Phosphorylates TERT at 'Ser-457', promoting TERT ubiquitination by the EDVP complex. Phosphorylates SIAH2, and thereby increases its ubiquitin ligase activity. Promotes the proteasomal degradation of MYC and JUN, and thereby regulates progress through the mitotic cell cycle and cell proliferation. Promotes proteasomal degradation of GLI2 and GLI3, and thereby plays a role in smoothened and sonic hedgehog signaling. Plays a role in cytoskeleton organization and neurite outgrowth via its phosphorylation of DCX and DPYSL2. Phosphorylates CRMP2/DPYSL2, CRMP4/DPYSL3, DCX, EIF2B5, EIF4EBP1, GLI2, GLI3, GYS1, JUN, MDM2, MYC, NFATC1, p53/TP53, TAU/MAPT and KATNA1. Can phosphorylate histone H1, histone H3 and histone H2B (in vitro). Can phosphorylate CARHSP1 (in vitro).
Cellular Location	Cytoplasm. Nucleus. Note=Translocates into the nucleus following DNA damage
Tissue Location	Testis, after the onset of spermatogenesis.

References

Becker W.,et al.J. Biol. Chem. 273:25893-25902(1998). Ota T.,et al.Nat. Genet. 36:40-45(2004). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases. Becker W.,et al.Submitted (NOV-1996) to the EMBL/GenBank/DDBJ databases. Woods Y.L.,et al.Biochem. J. 355:609-615(2001).

Images



WB Suggested Anti-DYRK2 Antibody Titration: 1.0 $\mu g/ml$ Positive Control: Fetal Lung

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.