

SNX2 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15119

Product Information

Application	WB
Primary Accession	O60749
Other Accession	NM_003100 , NP_003091
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58471

Additional Information

Gene ID	6643
Alias Symbol	MGC5204, TRG-9
Other Names	Sorting nexin-2, Transformation-related gene 9 protein, TRG-9, SNX2
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-SNX2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SNX2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SNX2
Function	Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2) (PubMed: 16179610). Acts in part as component of the retromer membrane- deforming SNX-BAR subcomplex (PubMed: 17101778). The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Can sense membrane curvature and has in vitro vesicle-to-membrane remodeling activity (PubMed: 23085988). Required for

retrograde endosome-to-TGN transport of TGN38 (PubMed:[20138391](#)). Promotes KALRN- and RHOG-dependent but retromer-independent membrane remodeling such as lamellipodium formation; the function is dependent on GEF activity of KALRN (PubMed:[20604901](#)).


Cellular Location

Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium Note=Colocalized with SORT1 to tubular endosomal membrane structures called endosome-to-TGN transport carriers (ETCs) which are budding from early endosome vacuoles just before maturing into late endosome vacuoles (PubMed:18088323). Colocalized with F-actin at the leading edge of lamellipodia in cells in a KALRN-dependent manner (PubMed:20604901).

References

Kurten R.C.,et al.Submitted (JAN-1998) to the EMBL/GenBank/DDBJ databases.
Haft C.R.,et al.Mol. Cell. Biol. 18:7278-7287(1998).
Kim J.W.,et al.Submitted (APR-2003) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



WB Suggested Anti-SNX2 Antibody Titration: 1.0 µg/ml
Positive Control: Hela Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.