

# USO1 antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15125

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">O60763</a>
<b>Other Accession</b>	<a href="#">NM_003715</a> , <a href="#">NP_003706</a>
<b>Reactivity</b>	Human, Rat, Pig, Dog, Horse
<b>Predicted</b>	Human, Rat, Pig, Dog, Horse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	107895

## Additional Information

---

<b>Gene ID</b>	8615
<b>Alias Symbol</b>	P115, TAP, VDP
<b>Other Names</b>	General vesicular transport factor p115, Protein USO1 homolog, Transcytosis-associated protein, TAP, Vesicle-docking protein, USO1, VDP
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 50 ul of distilled water. Final anti-USO1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	USO1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	USO1
<b>Synonyms</b>	VDP
<b>Function</b>	General vesicular transport factor required for intercisternal transport in the Golgi stack; it is required for transcytotic fusion and/or subsequent binding of the vesicles to the target membrane. May well act as a vesicular anchor by interacting with the target membrane and holding the vesicular and target membranes in proximity.
<b>Cellular Location</b>	Cytoplasm, cytosol. Golgi apparatus membrane; Peripheral membrane protein. Note=Recycles between the cytosol and the Golgi apparatus during

interphase. During interphase, the phosphorylated form is found exclusively in cytosol; the unphosphorylated form is associated with Golgi apparatus membranes


## References

---

Sohda M.,et al.J. Biol. Chem. 273:5385-5388(1998).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Bechtel S.,et al.BMC Genomics 8:399-399(2007).  
Hillier L.W.,et al.Nature 434:724-731(2005).  
Olsen J.V.,et al.Cell 127:635-648(2006).

## Images

---



WB Suggested Anti-USO1 Antibody Titration: 1.0 µg/ml  
Positive Control: 293T Whole CellUSO1 is strongly supported by BioGPS gene expression data to be expressed in Human HEK293T cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.