

TCAM2 Antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15166

Product Information

Application	WB
Primary Accession	Q86XR7
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26916

Additional Information

Gene ID	100302736;353376
Alias Symbol Other Names	TICAM2, TIRAP3, TIRP, TRAM, TIR domain-containing adapter molecule 2, TICAM-2, Putative NF-kappa-B-activating protein 502, TRIF-related adapter molecule, Toll-like receptor adaptor protein 3, Toll/interleukin-1 receptor domain-containing protein, MyD88-4, TICAM2, TIRAP3, TIRP, TRAM
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 μ l of distilled water. Final Anti-TCAM2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	TCAM2 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TICAM2
Synonyms	TIRAP3, TIRP, TRAM
Function	Functions as a sorting adapter in different signaling pathways to facilitate downstream signaling leading to type I interferon induction (PubMed: 16603631 , PubMed: 16757566 , PubMed: 25385819 , PubMed: 25825441). In TLR4 signaling, physically bridges TLR4 and TICAM1 and functionally transmits signal to TICAM1 in early endosomes after endocytosis of TLR4. In TLR2 signaling, physically bridges TLR2 and MYD88 and is required for the TLR2- dependent movement of MYD88 to endosomes

following ligand engagement (PubMed:[25385819](#)). Involved in IL-18 signaling and is proposed to function as a sorting adapter for MYD88 in IL-18 signaling during adaptive immune response (PubMed:[22685567](#)). Forms a complex with RAB11FIP2 that is recruited to the phagosomes to promote the activation of the actin-regulatory GTPases RAC1 and CDC42 and subsequent phagocytosis of Gram-negative bacteria (PubMed:[30883606](#)).

Cellular Location

[Isoform 1]: Cytoplasm. Golgi apparatus. Cell membrane. Endoplasmic reticulum. Early endosome membrane. Late endosome membrane. Cell projection, phagocytic cup. Note=Localized to the plasma membrane as a result of myristoylation. Phosphorylation on Ser-16 leads to its depletion from the membrane. Upon LPS stimulation colocalizes with isoform 2 in late endosomes


Tissue Location

Expressed in spleen, prostate, testis, uterus, small intestine, colon, peripheral blood leukocytes, heart, placenta, lung, liver, skeletal muscle, and pancreas
Isoform 2 is ubiquitously expressed (at lower levels than isoform 1)

References

Bin L.-H.,et al.J. Biol. Chem. 278:24526-24532(2003).
Oshiumi H.,et al.J. Biol. Chem. 278:49751-49762(2003).
Fitzgerald K.A.,et al.J. Exp. Med. 198:1043-1055(2003).
Fitzgerald K.A.,et al.J. Exp. Med. 198:1451-1451(2003).
Nakajima T.,et al.Immunogenetics 60:727-735(2008).

Images



Host: Rabbit
Target Name: TCAM2
Sample Tissue: Fetal Lung Lysate
Antibody Dilution: 1.0µg/ml

Host: Rabbit
Target Name: TCAM2
Sample Tissue: Fetal Lung lysates
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.