

SLC22A9 antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI15192

Product Information

Application WB
Primary Accession Q8IVM8

Other Accession <u>NM 080866</u>, <u>NP 543142</u>

Reactivity Human
Predicted Human
Host Rabbit
Clonality Polyclonal
Calculated MW 62169

Additional Information

Gene ID 114571

Alias Symbol FLJ23666, HOAT4, OAT4, OAT7, UST3H, ust3

Other Names Solute carrier family 22 member 9, Organic anion transporter 7, SLC22A9,

hOAT4, OAT7, UST3

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 50 ul of distilled water. Final anti-SLC22A9 antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions SLC22A9 antibody - C-terminal region is for research use only and not for use

in diagnostic or therapeutic procedures.

Protein Information

Name SLC22A9

Synonyms hOAT4, OAT7, UST3

Function Sodium-independent organic anion transporter, exhibits high specificity for

sulfated conjugates of xenobiotics and steroid hormones such as estrone

3-sulfate (E1S) and dehydroepiandrosterone sulfate (DHEAS)

(PubMed:<u>17393504</u>, PubMed:<u>26239079</u>, PubMed:<u>28945155</u>). Can transport

the statin pravastatin and may contribute to its disposition into the

hepatocytes when the function of OATPs is compromised

(PubMed:<u>26239079</u>). It is specifically activated by 3 to 5 carbons- containing short-chain fatty acids/SCFAs, including propionate (propanoate), butyrate

(butanoate) and valerate (pentanoate) (PubMed:<u>17393504</u>). May operate the exchange of sulfated organic components against short-chain fatty acids/SCFAs, in particular butanoate, at the sinusoidal membrane of

hepatocytes (PubMed: 17393504).

Cellular LocationBasolateral cell membrane; Multi-pass membrane protein. Note=Enriched at


the sinusoidal part of the plasma membrane

Tissue Location Specifically expressed in liver (also at protein level).

References

Shin H.J., et al. Hepatology 45:1046-1055(2007). Engel K., et al. Submitted (NOV-2000) to the EMBL/GenBank/DDBJ databases. Takanaga H., et al. Submitted (MAY-2001) to the EMBL/GenBank/DDBJ databases. Guo J.H., et al. Submitted (OCT-2001) to the EMBL/GenBank/DDBJ databases. Ota T., et al. Nat. Genet. 36:40-45(2004).

Images



WB Suggested Anti-SLC22A9 Antibody Titration: 1.0 µg/ml Positive Control: Fetal Heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.