

EIF3J antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15212

Product Information

Application	WB
Primary Accession	O75822
Other Accession	NM_003758 , NP_003749
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29062

Additional Information

Gene ID	8669
Alias Symbol	EIF3S1, eIF3-alpha, eIF3-p35
Other Names	Eukaryotic translation initiation factor 3 subunit J {ECO:0000255 HAMAP-Rule:MF_03009}, eIF3j {ECO:0000255 HAMAP-Rule:MF_03009}, Eukaryotic translation initiation factor 3 subunit 1 {ECO:0000255 HAMAP-Rule:MF_03009}, eIF-3-alpha {ECO:0000255 HAMAP-Rule:MF_03009}, eIF3 p35 {ECO:0000255 HAMAP-Rule:MF_03009}, EIF3J {ECO:0000255 HAMAP-Rule:MF_03009}
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-EIF3J antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	EIF3J antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EIF3J {ECO:0000255 HAMAP-Rule:MF_03009}
Function	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: 25849773 , PubMed: 27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex

(43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:[25849773](#)).


Cellular Location

Cytoplasm {ECO:0000255 | HAMAP-Rule:MF_03009}.

References

Block K.L.,et al.J. Biol. Chem. 273:31901-31908(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Li W.B.,et al.Submitted (JUL-2004) to the EMBL/GenBank/DDBJ databases.
Zody M.C.,et al.Nature 440:671-675(2006).
Bienvenut W.V.,et al.Submitted (MAR-2008) to UniProtKB.

Images



Host: Rabbit
Target Name: EIF3J
Antibody Dilution: 1.0µg/ml
Sample Tissue: Jurkat cell lysate
EIF3J is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.