

CXXC4 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15242

Product Information

Application	WB
Primary Accession	Q9H2H0
Other Accession	NM_025212 , NP_079488
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20978

Additional Information

Alias Symbol	IDAX, MGC149872
Other Names	CXXC-type zinc finger protein 4, Inhibition of the Dvl and axin complex protein, CXXC4, IDAX
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-CXXC4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	CXXC4 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CXXC4
Synonyms	IDAX
Function	Acts as a negative regulator of the Wnt signaling pathway via its interaction with DVL1 (By similarity). Binds preferentially to DNA containing cytidine-phosphate-guanosine (CpG) dinucleotides over CpH (H=A, T, and C), hemimethylated-CpG and hemimethylated-hydroxymethyl- CpG (PubMed: 29276034).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q9EQC9}.


References

Hino S.,et al.Mol. Cell. Biol. 21:330-342(2001).

Hillier L.W.,et al.Nature 434:724-731(2005).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-CXXC4 Antibody Titration: 1.0 µg/ml
Positive Control: MCF7 Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.