

HOOK3 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15256

Product Information

Application	WB
Primary Accession	Q86VS8
Other Accession	NM_032410 , NP_115786
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Goat, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Goat, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83126

Additional Information

Gene ID	84376
Alias Symbol	FLJ31058, HK3
Other Names	Protein Hook homolog 3, h-hook3, hHK3, HOOK3
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-HOOK3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	HOOK3 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HOOK3 (HGNC:23576)
Function	Acts as an adapter protein linking the dynein motor complex to various cargos and converts dynein from a non-processive to a highly processive motor in the presence of dynactin. Facilitates the interaction between dynein and dynactin and activates dynein processivity (the ability to move along a microtubule for a long distance without falling off the track). Predominantly recruits 2 dyneins, which increases both the force and speed of the microtubule motor (PubMed: 25035494 , PubMed: 33734450). Component of the FTS/Hook/FHIP complex (FHF complex). The FHF complex may function to promote vesicle trafficking and/or fusion via the homotypic vesicular protein sorting complex (the HOPS complex). May regulate clearance of endocytosed

receptors such as MSR1. Participates in defining the architecture and localization of the Golgi complex. FHF complex promotes the distribution of AP-4 complex to the perinuclear area of the cell (PubMed:[32073997](#)).


Cellular Location

Cytoplasm, cytoskeleton. Golgi apparatus. Note=Enriched at the cis-face of the Golgi complex. Localizes to microtubule asters in prophase (PubMed:11238449). Localizes to the manchette in elongating spermatids (By similarity). {ECO:0000250|UniProtKB:Q8BUK6, ECO:0000269|PubMed:11238449}

References

Walenta J.H.,et al.J. Cell Biol. 152:923-934(2001).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Beausoleil S.A.,et al.Nat. Biotechnol. 24:1285-1292(2006).
Sano H.,et al.J. Biol. Chem. 282:7973-7981(2007).

Images



WB Suggested Anti-HOOK3 Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.