

OST4 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15258

Product Information

| | |
|--------------------------|---|
| Application | WB |
| Primary Accession | P0C6T2 |
| Other Accession | NM_001134693 , NP_001128165 |
| Reactivity | Human, Mouse, Rat, Zebrafish, Dog |
| Predicted | Human, Mouse, Rat, Zebrafish, Dog |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 4193 |

Additional Information

| | |
|-------------------------------------|---|
| Gene ID | 100128731 |
| Alias Symbol | DKFZp586A0722 |
| Other Names | Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 4, OST4 |
| Format | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. |
| Reconstitution & Storage | Add 50 ul of distilled water. Final anti-OST4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles. |
| Precautions | OST4 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|--|
| Name | OST4 (HGNC:32483) |
| Function | Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol- pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation (PubMed: 31831667). N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity. Specifically involved in maintaining stability of STT3A-containing OST complexes. |

Cellular Location


Endoplasmic reticulum. Endoplasmic reticulum membrane; Single- pass type III membrane protein. Note=The single transmembrane helix has a kink in the middle of the transmembrane span

References

Hillier L.W.,et al.Nature 434:724-731(2005).

Gayen S.,et al.Biochem. Biophys. Res. Commun. 409:572-576(2011).

Images



WB Suggested Anti-OST4 Antibody Titration: 1.0 µg/ml
Positive Control: MCF7 Whole Cell

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.