

CCDC99 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15481

Product Information

Application	WB
Primary Accession	Q96EA4
Other Accession	NM_017785 , NP_060255
Reactivity	Human, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Rat, Rabbit, Pig, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70172

Additional Information

Gene ID	54908
Alias Symbol Other Names	FLJ20364, FLJ40690, CCDC99 Protein Spindly {ECO:0000255 HAMAP-Rule:MF_03041}, hSpindly, Arsenite-related gene 1 protein, Coiled-coil domain-containing protein 99 {ECO:0000255 HAMAP-Rule:MF_03041}, Rhabdomyosarcoma antigen MU-RMS-40.4A, Spindle apparatus coiled-coil domain-containing protein 1 {ECO:0000255 HAMAP-Rule:MF_03041}, SPDL1 {ECO:0000255 HAMAP-Rule:MF_03041}
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-CCDC99 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	CCDC99 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SPDL1 {ECO:0000255 HAMAP-Rule:MF_03041}
Function	Required for the localization of dynein and dynactin to the mitotic kintochore. Dynein is believed to control the initial lateral interaction between the kinetochore and spindle microtubules and to facilitate the subsequent formation of end-on kinetochore-microtubule attachments mediated by the NDC80 complex. Also required for correct spindle orientation. Does not appear to be required for the removal of spindle assembly checkpoint (SAC)

proteins from the kinetochore upon bipolar spindle attachment (PubMed:[17576797](#), PubMed:[19468067](#)). Acts as an adapter protein linking the dynein motor complex to various cargos and converts dynein from a non-processive to a highly processive motor in the presence of dynactin. Facilitates the interaction between dynein and dynactin and activates dynein processivity (the ability to move along a microtubule for a long distance without falling off the track) (PubMed:[25035494](#)). Plays a role in cell migration (PubMed:[30258100](#)).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Nucleus Cytoplasm, cytoskeleton, spindle pole. Note=Localizes to the nucleus in interphase and to the kinetochore in early prometaphase. Relocalizes to the mitotic spindle pole before metaphase and is subsequently lost from the spindle poles after chromosome congression is completed. Removal of this protein from the kinetochore requires the dynein/dynactin complex

References

Gu Y.Q.,et al.Submitted (MAY-2000) to the EMBL/GenBank/DDBJ databases.
Behrends U.,et al.Submitted (JAN-2004) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Schmutz J.,et al.Nature 431:268-274(2004).
Griffis E.R.,et al.J. Cell Biol. 177:1005-1015(2007).

Images



Host: Rabbit
Target Name: CCDC99
Sample Tissue: Hela Whole cell lysate
S
Antibody Dilution: 1.0µg/mlSPDL1 is supported by BioGPS gene expression data to be expressed in HeLa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.