

# CCDC134 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15482

## Product Information

Application	WB
Primary Accession	<a href="#">Q9H6E4</a>
Other Accession	<a href="#">NM_024821</a> , <a href="#">NP_079097</a>
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26561

## Additional Information

Gene ID	79879
Alias Symbol	FLJ22349, MGC21013, dj821D11.3
Other Names	Coiled-coil domain-containing protein 134, CCDC134
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-CCDC134 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	CCDC134 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	CCDC134 {ECO:0000303 PubMed:39509507, ECO:0000312 HGNC:HGNC:26185}
Function	Molecular adapter required to prevent protein hyperglycosylation of HSP90B1: during translation, associates with nascent HSP90B1 and the STT3A catalytic component of the OST-A complex and tethers them to a specialized translocon that forms a microenvironment for HSP90B1 folding (PubMed: <a href="#">38670073</a> , PubMed: <a href="#">39509507</a> ). In the CCDC134-containing translocon, STT3A associates with the SRT pseudosubstrate motif of HSP90B1, preventing access to facultative glycosylation sites until folding is completed, preventing hyperglycosylation and subsequent degradation of HSP90B1 (PubMed: <a href="#">39509507</a> ). In extracellular secreted form, promotes proliferation and activation of CD8(+) T-cells, suggesting a cytokine- like function

(PubMed:[25125657](#)). May inhibit ERK and JNK signaling activity (PubMed:[18087676](#), PubMed:[23070808](#)). May suppress cell migration and invasion activity, via its effects on ERK and JNK signaling (PubMed:[23070808](#)). May also localize in the nucleus: enhances stability of the PCAF histone acetyltransferase (HAT) complex member TADA2A and thus promotes PCAF-mediated histone acetyltransferase activity (PubMed:[22644376](#)). Has a critical role in the regulation of osteogenesis and bone development (PubMed:[32181939](#)).

#### Cellular Location

Endoplasmic reticulum lumen. Secreted. Cytoplasm Nucleus. Note=Mainly localizes to the endoplasmic reticulum (PubMed:39509507). Accumulates in the nucleus in response to UV irradiation (PubMed:22644376)

#### Tissue Location

Expressed in cervical gland, cervical squamous epithelium, endometrium, stomach, kidney distal convoluted tubule, spermatogenic cells in testis, mammary gland, liver and striated muscle (at protein level) (PubMed:18087676, PubMed:23070808). Also detected in placenta (PubMed:18087676). Highest expression in testis relative to other tissues (PubMed:18087676). Detected in T cells and dendritic cells; highly expressed in activated CD8(+) T cells, and also expressed at lower levels in CD4(+) T cells (PubMed:25125657)


## References

---

Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Dunham I.,et al.Nature 402:489-495(1999).  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).

## Images

---



Host: Rabbit  
Target Name: CCDC134  
Sample Tissue: Fetal Kidney lysates  
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.