

# SAPCD2 Antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI15567

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q86UD0</a>
Other Accession	<a href="#">NM_178448</a> , <a href="#">NP_848543</a>
Reactivity	Human, Rabbit, Dog, Guinea Pig
Predicted	Human, Rabbit, Pig, Dog, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42637

## Additional Information

---

Gene ID	89958
Alias Symbol	C9orf140, p42.3
Other Names	Suppressor APC domain-containing protein 2, Tumor specificity and mitosis phase-dependent expression protein, TS/MDEP, p42.3, SAPCD2, C9orf140
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-SAPCD2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SAPCD2 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	SAPCD2 ( <a href="#">HGNC:28055</a> )
Synonyms	C9orf140
Function	Plays a role in planar mitotic spindle orientation in retinal progenitor cells (RPCs) and promotes the production of symmetric terminal divisions (By similarity). Negatively regulates the mitotic apical cortex localization of GPSM2 (PubMed: <a href="#">26766442</a> ). Involved also in positive regulation of cell proliferation and tumor cell growth (PubMed: <a href="#">23576022</a> , PubMed: <a href="#">23704824</a> ).
Cellular Location	Cytoplasm. Nucleus. Cytoplasm, cell cortex. Apical cell membrane {ECO:0000250 UniProtKB:Q9D818}. Cell junction, tight junction.

Note=Localized at the apical cortical region during the M phase. In horizontally retinal progenitor dividing cells, localized at the pole cortical region from prophase to telophase cells. In vertically retinal progenitor dividing cells, not detected at the pole cortical region at any stage of mitosis {ECO:0000250|UniProtKB:Q9D818}

## Tissue Location

Expressed in 5-month-old fetal tissues, including stomach, intestine, colon, liver, brain, lung, heart, spleen and kidney (PubMed:17525738). Undetectable in non-cancerous adult tissues (PubMed:17525738). Expressed in many primary gastric carcinoma, but almost not in adjacent normal mucosa (PubMed:17525738). Expressed preferentially in M and G1 phases, compared to S and G2 phases (PubMed:17525738). Expression is up-regulated in hepatocellular carcinoma (HCC) and colorectal cancer (CRC) tissues (at protein level) (PubMed:23704824).

## Images

---



Host: Rabbit  
Target Name: SAPCD2  
Sample Tissue: Fetal Lung lysates  
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.