

SDCCAG3 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15626

Product Information

Application	WB
Primary Accession	Q96C92
Other Accession	NM_006643 , NP_006634
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47961

Additional Information

Gene ID	10807
Alias Symbol	NY-CO-3
Other Names	Serologically defined colon cancer antigen 3, Antigen NY-CO-3, SDCCAG3
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-SDCCAG3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SDCCAG3 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ENTR1 (HGNC:10667)
Synonyms	SDCCAG3
Function	Endosome-associated protein that plays a role in membrane receptor sorting, cytokinesis and ciliogenesis (PubMed: 23108400 , PubMed: 25278552 , PubMed: 27767179). Involved in the endosome-to-plasma membrane trafficking and recycling of SNX27-retromer-dependent cargo proteins, such as GLUT1 (PubMed: 25278552). Involved in the regulation of cytokinesis; the function may involve PTPN13 and GIT1 (PubMed: 23108400). Plays a role in the formation of cilia (PubMed: 27767179). Involved in cargo protein localization, such as PKD2, at primary cilia (PubMed: 27767179). Involved in the presentation of the tumor necrosis factor (TNF) receptor TNFRSF1A on the

cell surface, and hence in the modulation of the TNF-induced apoptosis (By similarity).


Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:A2AIW0}. Early endosome. Endosome. Recycling endosome. Midbody Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body
Note=Colocalizes in a WASHC2-dependent manner with the retromer CSC complex at endosomes (PubMed:25278552). During cytokinesis colocalized with PTPN13 at the midbody (PubMed:23108400). Colocalizes with IFT88 and gamma-tubulin at the basal body of primary cilia (PubMed:27767179) Colocalizes with IFT88 and pericentrin at the centrosome (PubMed:27767179).

Tissue Location

Expressed in the colon (at protein level).

Images



Host: Rabbit
Target Name: SDCCAG3
Sample Tissue: Fetal Heart lysates
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.