

INTS10 Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15855

Product Information

Application	WB
Primary Accession	Q9NVR2
Other Accession	NM_018142 , NP_060612
Reactivity	Human, Mouse, Rat, Rabbit, Dog, Guinea Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	82236

Additional Information

Gene ID	55174
Alias Symbol	C8orf35, INT10
Other Names	Integrator complex subunit 10, Int10, INTS10, C8orf35
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-INTS10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	INTS10 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	INTS10 {ECO:0000303 PubMed:38823386, ECO:0000312 HGNC:HGNC:25548}
Function	Component of the integrator complex, a multiprotein complex that terminates RNA polymerase II (Pol II) transcription in the promoter-proximal region of genes (PubMed: 38570683 , PubMed: 38823386). The integrator complex provides a quality checkpoint during transcription elongation by driving premature transcription termination of transcripts that are unfavorably configured for transcriptional elongation: the complex terminates transcription by (1) catalyzing dephosphorylation of the C-terminal domain (CTD) of Pol II subunit POLR2A/RPB1 and SUPT5H/SPT5, (2) degrading the exiting nascent RNA transcript via endonuclease activity and (3) promoting the release of Pol II from bound DNA (PubMed: 38570683). The integrator

complex is also involved in terminating the synthesis of non-coding Pol II transcripts, such as enhancer RNAs (eRNAs), small nuclear RNAs (snRNAs), telomerase RNAs and long non-coding RNAs (lncRNAs) (PubMed:[16239144](#), PubMed:[32647223](#)). Within the integrator complex, INTS10 is part of the integrator tail module that acts as a platform for the recruitment of transcription factors at promoters (PubMed:[38823386](#)). May be not involved in the recruitment of cytoplasmic dynein to the nuclear envelope, probably as component of the integrator complex (PubMed:[23904267](#)).

Cellular Location Nucleus


Background

Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.

References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Baillat D.,et al.Cell 123:265-276(2005).
Daub H.,et al.Mol. Cell 31:438-448(2008).

Images



Host: Rabbit
Target Name: INTS10
Sample Tissue: ACHN Whole cell lysate
S
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.