

# KIAA0141 Antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI15861

## Product Information

Application	WB
Primary Accession	<a href="#">Q14154</a>
Other Accession	<a href="#">NM_014773</a> , <a href="#">NP_055588</a>
Reactivity	Human, Horse, Bovine
Predicted	Human, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55920

## Additional Information

Gene ID	9812
Alias Symbol	DELE
Other Names	Death ligand signal enhancer, KIAA0141, DELE
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-KIAA0141 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	KIAA0141 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

Name	DELE1 {ECO:0000303   PubMed:32132706, ECO:0000312   HGNC:HGNC:28969}
Function	Protein kinase activator that acts as a key activator of the integrated stress response (ISR) following various stresses, such as iron deficiency, mitochondrial stress or mitochondrial DNA breaks (PubMed: <a href="#">32132706</a> , PubMed: <a href="#">32132707</a> , PubMed: <a href="#">35388015</a> , PubMed: <a href="#">37327776</a> , PubMed: <a href="#">37550454</a> , PubMed: <a href="#">37832546</a> , PubMed: <a href="#">38340717</a> ). Detects impaired protein import and processing in mitochondria, activating the ISR (PubMed: <a href="#">35388015</a> ). May also required for the induction of death receptor-mediated apoptosis through the regulation of caspase activation (PubMed: <a href="#">20563667</a> ).
Cellular Location	[DAP3-binding cell death enhancer 1]: Mitochondrion. Mitochondrion outer

membrane. Mitochondrion inner membrane. Note=Imported in the mitochondrial matrix in absence of stress, leading to its degradation by LONP1 (PubMed:37327776). Localizes at the mitochondrial surface in response to iron deficiency: iron deficiency impairs mitochondrial import, promoting localization at the mitochondrial surface and stabilization (PubMed:37327776). Associates with the mitochondrion inner membrane in response to mitochondrial stress, leading to its proteolytic processing by OMA1, and generation of the AP3-binding cell death enhancer 1 short form (DELE1(S) or S-DELE1) (PubMed:32132707)

#### Tissue Location

Detected in liver, skeletal muscle, kidney, pancreas, spleen, thyroid, testis, ovary, small intestine and colon

## Background

---

Essential for the induction of death receptor-mediated apoptosis through the regulation of caspase activation.


## References

---

Nagase T.,et al.DNA Res. 2:167-174(1995).  
Harada T.,et al.Apoptosis 15:1247-1255(2010).

## Images

---



Host: Rabbit  
Target Name: KIAA0141  
Sample Tissue: HepG2 Whole cell lysate  
S  
Antibody Dilution: 1.0µg/ml KIAA0141 is supported by BioGPS gene expression data to be expressed in HepG2

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.