

# RNF2 Antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI16052

## Product Information

---

Application	WB
Primary Accession	<a href="#">Q99496</a>
Other Accession	<a href="#">XP_005245469</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Predicted	Human, Mouse, Rat, Rabbit, Pig, Dog, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37655

## Additional Information

---

Gene ID	6045
Alias Symbol Other Names	RNF2, BAP1, DING, HIPI3, RING1B, E3 ubiquitin-protein ligase RING2, 6.3.2.-, Huntingtin-interacting protein 2-interacting protein 3, HIP2-interacting protein 3, Protein DinG, RING finger protein 1B, RING1b, RING finger protein 2, RING finger protein BAP-1, RNF2, BAP1, DING, HIPI3, RING1B
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 &mu; l of distilled water. Final Anti-RNF2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	RNF2 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	RNF2
Synonyms	BAP1, DING, HIPI3, RING1B
Function	E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A (H2AK119Ub), thereby playing a central role in histone code and gene regulation (PubMed: <a href="#">15386022</a> , PubMed: <a href="#">16359901</a> , PubMed: <a href="#">21772249</a> , PubMed: <a href="#">25355358</a> , PubMed: <a href="#">25519132</a> , PubMed: <a href="#">26151332</a> , PubMed: <a href="#">33864376</a> ). H2AK119Ub gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of

female mammals. May be involved in the initiation of both imprinted and random X inactivation (By similarity). Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:[16359901](#), PubMed:[26151332](#)). PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility (PubMed:[26151332](#)). E3 ubiquitin-protein ligase activity is enhanced by BMI1/PCGF4 (PubMed:[21772249](#)). Acts as the main E3 ubiquitin ligase on histone H2A of the PRC1 complex, while RING1 may rather act as a modulator of RNF2/RING2 activity (Probable). Association with the chromosomal DNA is cell-cycle dependent. In resting B- and T-lymphocytes, interaction with AURKB leads to block its activity, thereby maintaining transcription in resting lymphocytes (By similarity). Also acts as a negative regulator of autophagy by mediating ubiquitination of AMBRA1, leading to its subsequent degradation (By similarity).

#### Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q9CQJ4}. Chromosome {ECO:0000250|UniProtKB:Q9CQJ4}. Note=Enriched on inactive X chromosome (Xi) in female trophoblast stem (TS) cells as well as differentiating embryonic stem (ES) cells. The enrichment on Xi is transient during TS and ES cell differentiation. The association with Xi is mitotically stable in non-differentiated TS cells. Co-localizes with SAMD7 in nuclear polycomb bodies. {ECO:0000250|UniProtKB:Q9CQJ4}

## Background

---

E3 ubiquitin-protein ligase that mediates monoubiquitination of 'Lys-119' of histone H2A (H2AK119Ub), thereby playing a central role in histone code and gene regulation. H2AK119Ub gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. May be involved in the initiation of both imprinted and random X inactivation. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. E3 ubiquitin-protein ligase activity is enhanced by BMI1/PCGF4. Acts as the main E3 ubiquitin ligase on histone H2A of the PRC1 complex, while RING1 may rather act as a modulator of RNF2/RING2 activity. In resting B- and T-lymphocytes, interaction with AURKB leads to block its activity, thereby maintaining transcription in resting lymphocytes.

## References


---

Dyer M.J.S.,et al.Submitted (JAN-1997) to the EMBL/GenBank/DDBJ databases.  
 Li S.-F.,et al.Submitted (APR-1999) to the EMBL/GenBank/DDBJ databases.  
 Ota T.,et al.Nat. Genet. 36:40-45(2004).  
 Gregory S.G.,et al.Nature 441:315-321(2006).  
 Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

## Images

---

Host: Rabbit  
 Target Name: RNF2  
 Sample Tissue: MDA-MB-435s Whole cell lysate  
 s  
 Antibody Dilution: 1.0µg/ml



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.