

PDE10A Antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI16056

Product Information

Application	WB
Primary Accession	Q9Y233
Reactivity	Human, Rat, Rabbit, Dog, Horse
Predicted	Human, Rat, Rabbit, Dog, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	114946

Additional Information

Gene ID	10846
Alias Symbol Other Names	PDE10A, cAMP and cAMP-inhibited cGMP 3', 5'-cyclic phosphodiesterase 10A, 3.1.4.17, 3.1.4.35, PDE10A
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 µl of distilled water. Final Anti-PDE10A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	PDE10A Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PDE10A
Function	Plays a role in signal transduction by regulating the intracellular concentration of cyclic nucleotides (PubMed: 10373451 , PubMed: 10393245 , PubMed: 16330539 , PubMed: 17389385 , PubMed: 27058447). Can hydrolyze both cAMP and cGMP, but has higher affinity for cAMP and is more efficient with cAMP as substrate (PubMed: 10373451 , PubMed: 10393245 , PubMed: 17389385 , PubMed: 27058447). May play a critical role in regulating cAMP and cGMP levels in the striatum, a region of the brain that contributes to the control of movement and cognition (PubMed: 27058447).
Cellular Location	Cytoplasm, cytosol

Tissue Location

Abundant in the putamen and caudate nucleus regions of brain and testis, moderately expressed in the thyroid gland, pituitary gland, thalamus and cerebellum.

Background

Plays a role in signal transduction by regulating the intracellular concentration of cyclic nucleotides. Can hydrolyze both cAMP and cGMP, but has higher affinity for cAMP and is more efficient with cAMP as substrate.

References

Kotera J.,et al.Biochem. Biophys. Res. Commun. 261:551-557(1999).


Fujishige K.,et al.J. Biol. Chem. 274:18438-18445(1999).

Loughney K.,et al.Gene 234:109-117(1999).

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Mungall A.J.,et al.Nature 425:805-811(2003).

Images



Host: Rabbit
Target Name: PDE10A
Sample Tissue: PANC1 Whole cell lysate
S
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.