

DFFB Antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI16100

Product Information

Application	WB
Primary Accession	O76075
Other Accession	NP_004393
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39110

Additional Information

Gene ID	1677
Alias Symbol Other Names	DFFB, CAD, DFF2, DFF40, DNA fragmentation factor subunit beta, 3.-.-., Caspase-activated deoxyribonuclease, CAD, Caspase-activated DNase, Caspase-activated nuclease, CPAN, DNA fragmentation factor 40 kDa subunit, DFF-40, DFFB, CAD, DFF2, DFF40
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 μ l of distilled water. Final Anti-DFFB antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.
Precautions	DFFB Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DFFB
Synonyms	CAD, DFF2, DFF40
Function	Nuclease that induces DNA fragmentation and chromatin condensation during apoptosis. Degrades naked DNA and induces apoptotic morphology.
Cellular Location	Cytoplasm. Nucleus.


Background

Nuclease that induces DNA fragmentation and chromatin condensation during apoptosis. Degrades naked DNA and induces apoptotic morphology.

References

Liu X.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:8461-8466(1998).
Mukae N.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:9123-9128(1998).
Halenbeck R.,et al.Curr. Biol. 8:537-540(1998).
Nakagawara A.,et al.Submitted (JUN-1999) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Host: Rabbit
Target Name: DFFB
Sample Tissue: Jurkat Whole Cell lysates
Antibody Dilution: 1.0µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.