

FOXO4 antibody - middle region

Rabbit Polyclonal Antibody

Catalog # AI16169

Product Information

Application	WB
Primary Accession	P98177
Other Accession	NM_005938 , NP_005929
Reactivity	Human, Mouse, Rat, Pig, Dog, Horse
Predicted	Human, Mouse, Rat, Pig, Dog, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53684

Additional Information

Gene ID	4303
Alias Symbol Other Names	AFX, AFX1, MGC120490, MLLT7 Forkhead box protein O4, Fork head domain transcription factor AFX1, FOXO4, AFX, AFX1, MLLT7
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-FOXO4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	FOXO4 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FOXO4
Synonyms	AFX, AFX1, MLLT7
Function	Transcription factor involved in the regulation of the insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGF1. Down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle. Involved in increased proteasome activity in embryonic stem cells (ESCs) by activating expression of PSMD11 in ESCs, leading to enhanced assembly of the 26S proteasome, followed by higher proteasome activity.

Cellular Location	Cytoplasm. Nucleus. Note=When phosphorylated, translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation. Monoubiquitination increases nuclear localization. When deubiquitinated, translocated from nucleus to cytoplasm
Tissue Location	Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform zeta is most abundant in the liver, kidney, and pancreas


Background

Transcription factor involved in the regulation of the insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGFBP1. Down-regulates expression of HIF1A and suppresses hypoxia-induced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle. Involved in increased proteasome activity in embryonic stem cells (ESCs) by activating expression of PSMD11 in ESCs, leading to enhanced assembly of the 26S proteasome, followed by higher proteasome activity.

References

Peters U.,et al.Hum. Genet. 100:569-572(1997).
 Borkhardt A.,et al.Oncogene 14:195-202(1997).
 Yang Z.,et al.J. Biol. Chem. 277:8068-8075(2002).
 Ross M.T.,et al.Nature 434:325-337(2005).
 Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



WB Suggested Anti-FOXO4 Antibody Titration: 0.2-1 µg/ml
 ELISA Titer: 1:2500
 Positive Control: 293T cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.