

# HTR7 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI16211

#### **Product Information**

Application WB Primary Accession P34969

Other Accession <u>NM 019859</u>, <u>NP 062873</u>

**Reactivity Predicted**Human, Mouse, Rat, Pig, Dog, Horse
Human, Mouse, Rat, Pig, Dog, Horse

HostRabbitClonalityPolyclonalCalculated MW53555

#### **Additional Information**

Gene ID 3363

Alias Symbol 5-HT7

Other Names 5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT-X, Serotonin receptor 7,

HIR/

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

**Reconstitution & Storage** Add 50 ul of distilled water. Final anti-HTR7 antibody concentration is 1 mg/ml

in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C.

Avoid repeat freeze-thaw cycles.

**Precautions** HTR7 antibody - N-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name HTR7 {ECO:0000303 | Ref.3, ECO:0000312 | HGNC:HGNC:5302}

**Function** G-protein coupled receptor for 5-hydroxytryptamine (serotonin), a biogenic

hormone that functions as a neurotransmitter, a hormone and a mitogen (PubMed:<u>35714614</u>, PubMed:<u>8226867</u>). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G

proteins) and modulates the activity of downstream effectors

(PubMed:35714614, PubMed:8226867). HTR7 is coupled to G(s) G alpha

proteins and mediates activation of adenylate cyclase activity

(PubMed: 35714614).

**Cellular Location** Cell membrane; Multi-pass membrane protein

[Isoform A]: Predominant isoform in spleen, caudate and hippocampus.

[Isoform D]: Minor isoform in terms of expression.


## **Background**

This is one of the several different receptors for 5- hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of this receptor is mediated by G proteins that stimulate adenylate cyclase.

#### References

Heidmann D.E.A.,et al.J. Neurochem. 68:1372-1381(1997). Bard J.A.,et al.J. Biol. Chem. 268:23422-23426(1993). King M.M.,et al.Submitted (DEC-2003) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004). Goshima N.,et al.Nat. Methods 5:1011-1017(2008).

### **Images**



WB Suggested Anti-HTR7 Antibody Titration: 0.125µg/ml

ELISA Titer: 1:1562500

Positive Control: Jurkat cell lysate

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.