

HTR7 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI16212

Product Information

Application	WB
Primary Accession	P34969
Other Accession	NM_000872 , NP_000863
Reactivity	Human, Mouse, Rat, Pig, Dog, Horse
Predicted	Human, Mouse, Rat, Pig, Dog, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53555

Additional Information

Gene ID	3363
Alias Symbol	5-HT7
Other Names	5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT7, 5-HT-X, Serotonin receptor 7, HTR7
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-HTR7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	HTR7 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	HTR7 {ECO:0000303 Ref.3, ECO:0000312 HGNC:HGNC:5302}
Function	G-protein coupled receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone and a mitogen (PubMed: 35714614 , PubMed: 8226867). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors (PubMed: 35714614 , PubMed: 8226867). HTR7 is coupled to G(s) G alpha proteins and mediates activation of adenylate cyclase activity (PubMed: 35714614).
Cellular Location	Cell membrane; Multi-pass membrane protein

Tissue Location

[Isoform A]: Predominant isoform in spleen, caudate and hippocampus.
[Isoform D]: Minor isoform in terms of expression.

Images



Host: Rabbit
Target Name: HTR7
Lane A: Marker
Lane B: Jurkat cell lysate
S
Antibody Dilution: 0.125 µg/ml

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.