

# ATF5 antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # AI16226

# **Product Information**

Application	WB, IHC
Primary Accession	<u>Q9Y2D1</u>
Other Accession	<u>NM_012068, NP_036200</u>
Reactivity	Human, Mouse, Rat, Pig, Guinea Pig, Bovine
Predicted	Human, Mouse, Rat, Pig, Guinea Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30674

# **Additional Information**

Gene ID	22809
Alias Symbol Other Names	ATFX, FLJ34666, HMFN0395 Cyclic AMP-dependent transcription factor ATF-5, cAMP-dependent transcription factor ATF-5, Activating transcription factor 5, Transcription factor ATFx, ATF5, ATFX
Format	Lyophilized
Reconstitution & Storage	Add 50 ul of distilled water. Final anti-ATF5 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	ATF5 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

### **Protein Information**

Name	ATF5
Synonyms	ATFX
Function	Transcription factor that either stimulates or represses gene transcription through binding of different DNA regulatory elements such as cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), ATF5-specific response element (ARE) (consensus: 5'- C[CT]TCT[CT]CCTT[AT]-3') but also the amino acid response element (AARE), present in many viral and cellular promoters. Critically involved, often in a cell type-dependent manner, in cell survival, proliferation, and differentiation (PubMed:10373550, PubMed:15358120, PubMed:20654631, PubMed:21212266). Its transcriptional activity is

	enhanced by CCND3 and slightly inhibited by CDK4 (PubMed: <u>15358120</u> ). Important regulator of the cerebral cortex formation, functions in cerebral cortical neuroprogenitor cells to maintain proliferation and to block differentiation into neurons. Must be down-regulated in order for such cells to exit the cycle and differentiate (By similarity). Participates in the pathways by which SHH promotes cerebellar granule neuron progenitor cells proliferation (By similarity). Critical for survival of mature olfactory sensory neurons (OSN), directs expression of OSN-specific genes (By similarity). May be involved in osteogenic differentiation (PubMed: <u>22442021</u> ). Promotes cell proliferation and survival by inducing the expression of EGR1 sinergistically with ELK1. Once acetylated by EP300, binds to ARE sequences on target genes promoters, such as BCL2 and EGR1 (PubMed: <u>21791614</u> ). Plays an anti-apoptotic role through the transcriptional regulation of BCL2, this function seems to be cell type-dependent (By similarity). Cooperates with NR113/CAR in the transcriptional activation of CYP2B6 in liver (PubMed: <u>18332083</u> ). In hepatic cells, represses CRE-dependent transcription and inhibits proliferation by blocking at G2/M phase (PubMed: <u>18701499</u> , PubMed: <u>22528486</u> ). May act as a negative regulator of IL1B transduction pathway in liver (PubMed: <u>24379400</u> ). Upon IL1B stimulus, cooperates with NLK to activate the transactivation activity of C/EBP subfamily members (PubMed: <u>24216764</u> ). Regulates centrosome dynamics in a cell-cycle- and centriole-age-dependent manner. Forms 9-foci symmetrical ring scaffold around the mother centriole to control centrosome function and the interaction between centrioles and pericentriolar material (PubMed: <u>26213385</u> ).
Cellular Location	Cytoplasm. Nucleus {ECO:0000255 PROSITE-ProRule:PRU00978, ECO:0000269 PubMed:15358120, ECO:0000269 PubMed:22528486}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Actively transported to the centrosome and accumulated in the pericentriolar material (PCM) during G1 to M phase via a microtubule- dependent mechanism. During late telophase and cytokinesis, translocates from the centrosome to the midbody
Tissue Location	Widely expressed with higher expression levels in liver.

# Background

Transcriptional activator which binds the cAMP response element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters and blocks the differentiation of neuroprogenitor cells into neurons. Its transcriptional activity is enhanced by CCND3 and slightly inhibited by CDK4.

#### References

White J.H., et al. Proc. Natl. Acad. Sci. U.S.A. 97:13967-13972(2000). Yamada S., et al.Oncogene 23:5901-5911(2004). Kohroki J., et al.Submitted (DEC-1998) to the EMBL/GenBank/DDBJ databases. Ota T., et al.Nat. Genet. 36:40-45(2004). Grimwood J., et al.Nature 428:529-535(2004).

#### Images



WB Suggested Anti-ATF5 Antibody Titration: 0.2-1  $\mu$ g/ml ELISA Titer: 1:312500 Positive Control: Jurkat cell lysate ATF5 is strongly supported by BioGPS gene expression data to be expressed in Human Jurkat cells



Liver

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.