

SOX9 antibody - N-terminal region

Rabbit Polyclonal Antibody

Catalog # AI16242

Product Information

Application	WB
Primary Accession	P48436
Other Accession	NM_000346 , NP_000337
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Dog, Guinea Pig, Bovine, Neisseria Gonorrhoeae
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Dog, Guinea Pig, Bovine, Neisseria Gonorrhoeae
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56137

Additional Information

Gene ID	6662
Alias Symbol	CMD1, SRA1, CMPD1
Other Names	Transcription factor SOX-9, SOX9
Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Reconstitution & Storage	Add 100 ul of distilled water. Final anti-SOX9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
Precautions	SOX9 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SOX9 {ECO:0000303 PubMed:7990924, ECO:0000312 HGNC:HGNC:11204}
Function	Transcription factor that plays a key role in chondrocytes differentiation and skeletal development (PubMed: 24038782). Specifically binds the 5'-ACAAAG-3' DNA motif present in enhancers and super-enhancers and promotes expression of genes important for chondrogenesis, including cartilage matrix protein-coding genes COL2A1, COL4A2, COL9A1, COL11A2 and ACAN, SOX5 and SOX6 (PubMed: 8640233). Also binds to some promoter regions (By similarity). Plays a central role in successive steps of chondrocyte differentiation (By similarity). Absolutely required for precartilaginous condensation, the first step in chondrogenesis during which skeletal progenitors differentiate into prechondrocytes (By similarity). Together with

SOX5 and SOX6, required for overt chondrogenesis when condensed prechondrocytes differentiate into early stage chondrocytes, the second step in chondrogenesis (By similarity). Later, required to direct hypertrophic maturation and block osteoblast differentiation of growth plate chondrocytes: maintains chondrocyte columnar proliferation, delays prehypertrophy and then prevents osteoblastic differentiation of chondrocytes by lowering beta-catenin (CTNNB1) signaling and RUNX2 expression (By similarity). Also required for chondrocyte hypertrophy, both indirectly, by keeping the lineage fate of chondrocytes, and directly, by remaining present in upper hypertrophic cells and transactivating COL10A1 along with MEF2C (By similarity). Low lipid levels are the main nutritional determinant for chondrogenic commitment of skeletal progenitor cells: when lipids levels are low, FOXO (FOXO1 and FOXO3) transcription factors promote expression of SOX9, which induces chondrogenic commitment and suppresses fatty acid oxidation (By similarity). Mechanistically, helps, but is not required, to remove epigenetic signatures of transcriptional repression and deposit active promoter and enhancer marks at chondrocyte-specific genes (By similarity). Acts in cooperation with the Hedgehog pathway-dependent GLI (GLI1 and GLI3) transcription factors (By similarity). In addition to cartilage development, also acts as a regulator of proliferation and differentiation in epithelial stem/progenitor cells: involved in the lung epithelium during branching morphogenesis, by balancing proliferation and differentiation and regulating the extracellular matrix (By similarity). Controls epithelial branching during kidney development (By similarity).

Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00267, ECO:0000269 | PubMed:8640233}


Background

Plays an important role in the normal skeletal development. May regulate the expression of other genes involved in chondrogenesis by acting as a transcription factor for these genes.

References

Foster J.W.,et al.Nature 372:525-530(1994).
Wagner T.,et al.Cell 79:1111-1120(1994).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Cox J.J.,et al.N. Engl. J. Med. 364:91-93(2011).

Images



WB Suggested Anti-SOX9 Antibody Titration: 2.5µg/ml
ELISA Titer: 1:12500
Positive Control: HepG2 cell lysate
SOX9 is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells