

GABPA antibody - C-terminal region

Rabbit Polyclonal Antibody Catalog # AI16246

Product Information

Application WB, IHC Primary Accession Q06546

Other Accession NM 002040, NP 002031

Reactivity Human, Mouse, Rat, Dog, Horse, Bovine

Predicted Human, Mouse, Rat, Chicken, Dog, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 51295

Additional Information

Gene ID 2551

Alias Symbol NFT2, NRF2, NRF2A, E4TF1A, E4TF1-60, RCH04A07

Other Names GA-binding protein alpha chain, GABP subunit alpha, Nuclear respiratory

factor 2 subunit alpha, Transcription factor E4TF1-60, GABPA, E4TF1A

Format Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium

azide and 2% sucrose.

Reconstitution & Storage Add 100 ul of distilled water. Final anti-GABPA antibody concentration is 1

mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at

20°C. Avoid repeat freeze-thaw cycles.

Precautions GABPA antibody - C-terminal region is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name GABPA

Synonyms E4TF1A


Function Transcription factor capable of interacting with purine rich repeats (GA

repeats). Positively regulates transcription of transcriptional repressor


RHIT/ZNF205 (PubMed:22306510).

Cellular Location Nucleus.

Images



Human kidney



WB Suggested Anti-GABPA Antibody Titration: 0.5-1.0µg/ml
Positive Control: Jurkat cell lysate
GABPA is supported by BioGPS gene expression data to be expressed in Jurkat

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.