

# GABPA antibody - C-terminal region

Rabbit Polyclonal Antibody

Catalog # AI16246

## Product Information

---

<b>Application</b>	WB, IHC
<b>Primary Accession</b>	<a href="#">Q06546</a>
<b>Other Accession</b>	<a href="#">NM_002040</a> , <a href="#">NP_002031</a>
<b>Reactivity</b>	Human, Mouse, Rat, Dog, Horse, Bovine
<b>Predicted</b>	Human, Mouse, Rat, Chicken, Dog, Horse, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	51295

## Additional Information

---

<b>Gene ID</b>	2551
<b>Alias Symbol</b>	NFT2, NRF2, NRF2A, E4TF1A, E4TF1-60, RCH04A07
<b>Other Names</b>	GA-binding protein alpha chain, GABP subunit alpha, Nuclear respiratory factor 2 subunit alpha, Transcription factor E4TF1-60, GABPA, E4TF1A
<b>Format</b>	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
<b>Reconstitution &amp; Storage</b>	Add 100 ul of distilled water. Final anti-GABPA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.
<b>Precautions</b>	GABPA antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


## Protein Information

---

<b>Name</b>	GABPA
<b>Synonyms</b>	E4TF1A
<b>Function</b>	Transcription factor capable of interacting with purine rich repeats (GA repeats). Positively regulates transcription of transcriptional repressor RHIT/ZNF205 (PubMed: <a href="#">22306510</a> ).
<b>Cellular Location</b>	Nucleus.

## Images

---



Human kidney



WB Suggested Anti-GABPA Antibody Titration:

0.5-1.0µg/ml

Positive Control: Jurkat cell lysate

GABPA is supported by BioGPS gene expression data to be expressed in Jurkat

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.