

Anti-SPI1 / PU.1 Antibody

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS17274

Product Information

Application	WB, IHC-P, E
Primary Accession	P17947
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	31083

Additional Information

Gene ID	6688
Alias Symbol	SPI1
Other Names	SPI1, 31 kDa-transforming protein, 31 kDa transforming protein, PU.1, SFPI1, SPI-1, SPI-1 proto-oncogene, SPI-A, Transcription factor PU.1
Target/Specificity	Human SPI1 / PU.1
Reconstitution & Storage	PBS, pH 7.4, 0.03% Proclin 300, 50% glycerol. Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.
Precautions	Anti-SPI1 / PU.1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SPI1
Function	Pioneer transcription factor, which controls hematopoietic cell fate by decompacting stem cell heterochromatin and allowing other transcription factors to enter otherwise inaccessible genomic sites. Once in open chromatin, can directly control gene expression by binding genetic regulatory elements and can also more broadly influence transcription by recruiting transcription factors, such as interferon regulatory factors (IRFs), to otherwise inaccessible genomic regions (PubMed: 23658224 , PubMed: 33951726). Transcriptionally activates genes important for myeloid and lymphoid lineages, such as CSF1R (By similarity). Transcriptional activation from certain promoters, possibly containing low affinity binding sites, is achieved cooperatively with other transcription factors. FCER1A transactivation is achieved in cooperation with GATA1 (By similarity). May be particularly important for the pro- to pre-B cell transition (PubMed: 33951726). Binds (via

the ETS domain) onto the purine-rich DNA core sequence 5'-GAGGAA-3', also known as the PU-box (PubMed:[33951726](#)). In vitro can bind RNA and interfere with pre-mRNA splicing (By similarity).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00237, ECO:0000269 | PubMed:33951726}

Tissue Location

In the bone marrow, concentrated in hematopoietic stem cell, lymphoid progenitor, myeloid lineage (granulocyte macrophage progenitors, classical dendritic cells, monocytes) and B-cell clusters Among B-cells, predominantly expressed in pre-B1 cells (PubMed:33951726). Expressed in germinal center B-cells (PubMed:23166356).

Images



Human Tonsil: Formalin-Fixed, Paraffin-Embedded (FFPE)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.